


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1997

[Full Table of Content](#)



97/R/WW/8 Fungicides and Take-all - W. Wheat

Rothamsted Research

Rothamsted Research (1998) *97/R/WW/8 Fungicides and Take-all - W. Wheat ; Yields Of The Field Experiments 1997*, pp 117 - 118 - DOI: <https://doi.org/10.23637/ERADOC-1-53>

97/R/WW/8

WINTER WHEAT

FUNGICIDES AND TAKE-ALL

Object: To test fungicides to control take-all (*Gaeumannomyces graminis*) and eyespot (*Tapesia yellundae*) - Stubbings.

Sponsors: G.L. Bateman, R.J. Gutteridge.

Design: 4 randomised blocks of 6 x 2 plots.

Whole plot dimensions: 3.0 x 10.0.

Treatments:

- | | |
|-------------|--|
| 1. SEED TRT | Seed treatment: |
| - | None |
| F1R1 | Fungicide 1 rate 1 |
| F1R2 | Fungicide 1 rate 2 |
| F2R1 | Fungicide 2 rate 1 |
| F2R2 | Fungicide 2 rate 2 |
| B | Fuberidazole and triadimenol (Baytan Flowable) |
| 2. SPORTAK | Eyespot control: |
| O | None |
| S | Prochloraz (Sportak 45) |

NOTE: Fungicide 1 and 2 are under commercial development, composition undisclosed.

Experimental diary:

28-Aug-96 : B : PK as (0:20:32) at 1400 kg.
05-Oct-96 : B : Ploughed and furrow pressed.
14-Oct-96 : T : Rotary harrowed, Brigadier, dressed as treatment, drilled at 380 seeds per m².
28-Nov-96 : B : Auger at 2.5 l with Stomp 400 SC at 3.1 l in 200 l.
10-Mar-97 : B : 34.5% N at 118 kg.
01-Apr-97 : T : **SPORTAK** S: Sportak 45 at 0.9 l in 220 l.
07-Apr-97 : B : 34.5% N at 464 kg.
04-Jun-97 : B : Folicur at 0.5 l with Mallard 750 EC at 0.3 l and Pointer at 0.5 l in 300 l.
09-Jun-97 : B : Mistral at 1.0 l in 300 l.
12-Aug-97 : B : Combine harvested.

Previous crops: W. wheat 1995 and 1996.

NOTE: Plant samples were taken in April and July to assess take-all and stem base diseases, Septoria (*Septoria tritici*) was also assessed in July.

97/R/WW/8

GRAIN TONNES/HECTARE

***** Tables of means *****

SPORTAK SEED TRT	O	S	Mean
-	7.60	8.08	7.84
F1R1	8.63	8.70	8.67
F1R2	8.29	8.34	8.31
F2R1	8.05	7.88	7.97
F2R2	8.28	8.27	8.28
B	7.88	8.02	7.95
Mean	8.12	8.21	8.17

*** Standard errors of differences of means ***

SEED TRT	SPORTAK	SEED TRT SPORTAK
0.174	0.100	0.246

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	33	0.347	4.3
GRAIN MEAN DM%	86.5		
PLOT AREA HARVESTED	0.00237		