


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1997

[Full Table of Content](#)



97/R/RAS/3 Neem Study - S. Oilseed Rape

Rothamsted Research

Rothamsted Research (1998) *97/R/RAS/3 Neem Study - S. Oilseed Rape* ; Yields Of The Field Experiments 1997, pp 136 - 137 - DOI: <https://doi.org/10.23637/ERADOC-1-53>

97/R/RAS/3

SPRING OILSEED RAPE

NEEM STUDY

Object: To test two formulations of neem on cereal aphids in spring wheat - Delafield.

Sponsors: L.E. Smart, B.J. Pye.

Design: 4 randomised blocks of 6 plots.

Whole plot dimensions: 3.0 x 10.0.

Treatments:

NEEMFORM Formulation and timing:

-	None
I	Deltamethrin at aphid migration
O1	Neem oil at 2.5 kg in 10.4 l by electrostatic sprayer
O2	Neem oil at 2.5 kg in 200 l by hydraulic sprayer
E1	Neem extract at 7.5 g a.i. in 10.4 l by electrostatic sprayer
E2	Neem extract at 7.5 g a.i. in 200 l by hydraulic sprayer

Experimental diary:

31-Oct-96 : B : Ploughed.
21-Mar-97 : B : Rolled. Spring-tine cultivated.
06-Apr-97 : B : MTM Trifluralin at 2.3 l in 390 l. Spring-tine cultivated twice.
07-Apr-97 : B : Rotary harrowed, Starlight, recleaned, drilled at 120 seeds per m². Rolled.
09-Apr-97 : B : Irrigated 13 mm.
18-Apr-97 : B : Irrigated 13 mm.
01-May-97 : B : Cyperkill 10 at 300 ml with Vassgro Non-ionic at 50 ml in 300 l.
15-May-97 : B : Cyperkill 10 at 300 ml with Vassgro Non-ionic at 50 ml in 300 l.
30-May-97 : B : 34.5% N at 290 kg.
10-Jun-97 : T : **NEEMFORM** I: Decis at 0.5 l in 200 l.
 : T : **NEEMFORM** O1, O2, E1 and E2: Neem treatments applied.
17-Jun-97 : T : **NEEMFORM** O1, O2, E1 and E2: Neem treatments applied.
24-Jun-97 : T : **NEEMFORM** O1, O2, E1 and E2: Neem treatments applied.
04-Sep-97 : B : Combine harvested.

Previous crops: S. rape 1995, w. wheat 1996.

NOTES: (1) Neem products applied at 2.5 l product in 200 l for hydraulic sprayer and in 10.4 l for electrostatic sprayer.
(2) Pollen beetle populations were assessed weekly through the flowering period.

97/R/RAS/3

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

NEEMFORM

-	1.25
I	1.33
O1	1.39
O2	1.25
E1	1.39
E2	1.19

Mean	1.30
------	------

*** Standard errors of differences of means ***

NEEMFORM

0.101

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	15	0.143	11.0

GRAIN MEAN DM% 83.6

PLOT AREA HARVESTED 0.00240