

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1998

[Full Table of Content](#)



98/R/CS/494 Take-all, Phialophora and Seed Treatments - S. Wheat

Rothamsted Research

Rothamsted Research (1999) *98/R/CS/494 Take-all, Phialophora and Seed Treatments - S. Wheat* ; Yields Of The Field Experiments 1998, pp 114 - 115 - DOI:

<https://doi.org/10.23637/ERADOC-1-52>

98/R/CS/494

TAKE-ALL, PHIALOPHORA AND SEED TREATMENTS

Object: To test the effects in the field of a fungal antagonist of take-all (*Gaeumannomyces graminis*) and determine its interaction with a take-all-selective fungicide - Stackyard.

Sponsors: G.L. Bateman, R.J. Gutteridge.

The first year, s. wheat.

Design: 4 randomised blocks of 18 plots.

Plot dimensions: 3.0 x 10.0.

Treatments: Combinations of:-

1. **TRT1998** Treatment to 1998 crop:

-8	None
P8	<i>Phialophora</i> inoculum to seedbed
T8	Take-all inoculum to seedbed
2. **TRT1999** Treatment to 1999 crop:

-9	None
P9	<i>Phialophora</i> inoculum to seedbed
T9	Take-all inoculum to seedbed
3. **TRT2000** Treatment to 2000 crop:

-0	None
P0	<i>Phialophora</i> inoculum to seedbed
S0	Seed treatment fungicide
PS0	<i>Phialophora</i> and seed treatment

Experimental diary:

31-Oct-97 : B : Ploughed.
30-Jan-98 : B : Barclay Gallup at 2.0 l with Chiltern Cropoil at 1.0 l in 200 l.
12-Feb-98 : B : Parable at 2.0 l with Scythe LC at 0.5 l and Luxon Non-Ionic Wetter at 0.1 l in 200 l.
16-Mar-98 : B : Heavy spring-tine cultivated.
17-Mar-98 : B : Spring-tine cultivated.
18-Mar-98 : T : **TRT1998** P8: Inoculum applied at 20 g per m². Rotary harrowed.
 : T : **TRT1998** T8: Inoculum applied at 20 g per m². Rotary harrowed.
 : B : Rotary harrowed, Axona, undressed, drilled at 400 seeds per m².
29-Apr-98 : B : 34.5% N at 380 kg.
05-May-98 : B : Ally at 30 g with MSS Optica at 2.0 l in 200 l.
04-Jun-98 : B : Corbel at 0.3 l with Opus at 0.3 l in 200 l.
12-Jun-98 : B : Topik at 125 ml with Chiltern Cropoil at 1.0 l in 200 l.

98/R/CS/494

Experimental diary:

28-Jun-98 : B : Bavistin DF at 0.3 kg with Radar at 0.25 l in 200 l.
02-Sep-98 : B : Combine harvested.

NOTE: Plant samples were taken in July to assess take-all and root colonisation by *Phialophora*.

GRAIN TONNES/HECTARE

***** Tables of means *****

TRT1998

-8	7.02
P8	6.83
T8	5.81

Mean	6.63
------	------

*** Standard errors of differences of means ***

TRT1998

0.099	min.rep
0.089	max-min

TRT1998

min.rep	P8 v T8
max-min	-8 v P8 or T8

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	66	0.313	4.7

GRAIN MEAN DM% 82.0

PLOT AREA HARVESTED 0.00229