

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1998

[Full Table of Content](#)



98/R/LP/7 Genotype, Row Spacing and Seed Rate - Lupins

Rothamsted Research

Rothamsted Research (1999) *98/R/LP/7 Genotype, Row Spacing and Seed Rate - Lupins ; Yields Of The Field Experiments 1998*, pp 178 - 180 - DOI: <https://doi.org/10.23637/ERADOC-1-52>

98/R/LP/7

LUPINS

GENOTYPE, ROW SPACING AND SEED RATE

Object: To test seed rates and row spacings on the structure and yield of standard and new dwarf-determinate genotypes of lupins - Osier.

Sponsors: I.F. Shield, J.E. Leach.

Design: 3 blocks of (4 x 2 x 2) + 2 extra plots.

Whole plot dimensions: 8.0 x 9.0.

Treatments: All combinations of:-

1. GENOTYPE

LY	Lucyane dressed Germipro
LB	Ludet dressed Germipro
12	DTN 12 undressed
20	DTN 20 undressed

2. ROW SPAC

Row spacing (cm):

R1	12.5
R2	37.5

3. SEED RAT

Seed rate seeds per m²:

S1	40
S2	80

Plus extra

EXTRA

X1	Ludet at 50 cm and 40 seeds per m ²
X2	DTN 20 at 50 cm and 40 seeds per m ²

Experimental diary:

12-Aug-97 : B : Ploughed and furrow pressed.
15-Sep-97 : B : Rolled. Rotary harrowed.
16-Sep-97 : T : Genotypes drilled as treatment.
18-Sep-97 : B : Rolled. Stomp 400 SC at 5.0 l with Spannit at 1.1 l in 200 l.
27-Oct-97 : B : Checkmate at 1.75 l with Atlas Adjuvant Oil at 2.0 l in 200 l.
26-Nov-97 : B : Gesatop 500 SC at 2.5 l in 200 l.
18-Feb-98 : B : Folicur at 0.5 l in 200 l.
24-Feb-98 : B : Rovral Flo at 1.0 l in 200 l.
16-Mar-98 : B : Skirmish at 1.0 l in 200 l.
02-Apr-98 : B : Danadim Dimethoate 40 at 850 ml in 200 l.
20-May-98 : B : Folicur at 0.5 l in 200 l. Aphox at 280 g in 200 l.

98/R/LP/7

Experimental diary:

12-Jun-98 : B : Alto 240 EC at 0.33 l with Bravo 500 at 1.5 l in 200 l.
 17-Sep-98 : T : **GENOTYPE** LY, LB: Combine harvested.
 23-Sep-98 : T : **GENOTYPE** 12, 20: Combine harvested.

Previous crops: Set-aside and industrial rape 1996, w. wheat and w. barley 1997.

NOTE: Plant populations were counted after establishment and in April. Light interception was measured, plants were sampled to measure dry matter accumulation. Lodging was assessed before harvest and at harvest components of yield were measured.

GRAIN TONNES/HECTARE

***** Tables of means *****

ROW SPAC GENOTYPE	R1	R2	Mean
LY	1.91	2.22	2.07
LD	3.04	2.72	2.88
12	1.56	1.53	1.54
20	2.08	2.42	2.25
Mean	2.15	2.22	2.19

SEED RAT GENOTYPE	S1	S2	Mean
LY	1.99	2.14	2.07
LD	2.81	2.96	2.88
12	1.59	1.50	1.54
20	2.44	2.05	2.25
Mean	2.21	2.17	2.19

SEED RAT ROW SPAC	S1	S2	Mean
R1	2.23	2.06	2.15
R2	2.18	2.27	2.22
Mean	2.21	2.17	2.19

ROW SPAC SEED RAT GENOTYPE	R1 S1	R2 S2	R2 S1	S2
LY	1.84	1.99	2.14	2.30
LD	3.10	2.98	2.52	2.93
12	1.57	1.56	1.60	1.45
20	2.43	1.72	2.45	2.39

EXTRA	X1	X2	Mean
	1.93	0.66	1.30

Grand mean 2.09

98/R/LP/7

GRAIN TONNES/HECTARE

*** Standard errors of differences of means ***

GENOTYPE	ROW SPAC	SEED RAT	GENOTYPE
			ROW SPAC
0.173	0.122	0.122	0.244
GENOTYPE	ROW SPAC	GENOTYPE	
SEED RAT	SEED RAT	ROW SPAC	
		SEED RAT	
		& EXTRA	
0.224	0.173	0.345	

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	34	0.423	20.3

GRAIN MEAN DM% 75.1

PLOT AREA HARVESTED	ROW SPAC R1	0.00207
PLOT AREA HARVESTED	ROW SPAC R2	0.00202
PLOT AREA HARVESTED	EXTRA PLOTS	0.00225