

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1998

[Full Table of Content](#)



98/R/CS/457 Set-aside, Cultivations and Crops - W. Wheat

Rothamsted Research

Rothamsted Research (1999) *98/R/CS/457 Set-aside, Cultivations and Crops - W. Wheat ; Yields Of The Field Experiments 1998*, pp 90 - 92 - DOI: <https://doi.org/10.23637/ERADOC-1-52>

98/R/CS/457

SET-ASIDE, CULTIVATIONS AND CROPS

Object: To measure the establishment, growth and yield of w. wheat and w. rape following a range of cultivations and herbicide applications after natural regeneration set-aside. To assess soil nitrogen and weeds in the two crops and diseases in the wheat - Scout.

Sponsors: J.F. Jenkyn, R.G. Gutteridge.

The third year, w. wheat.

Design: 3 randomised blocks of 5 x 2 split into 2 sub-plots.

Whole plot dimensions: 12.0 x 26.0.

Sub-plot dimensions: 10.0 x 12.0.

Treatments: All combinations of:-

1. **SETDESTR** Method and time of destruction of set-aside in 1996:

 (PG) Ploughed in May, glyphosate pre-drilling
 (PC) Ploughed in May, cultivated in June and July
 (MP) Minimally cultivated in May, ploughed in August
 (HP) Herbicide in May, ploughed in August
 (-P) Ploughed in August
2. **CROP** Crop in 1997

 (R) Winter rape
 (W) Winter wheat

Sub-plots

3. **NITROGEN** Fertilizer nitrogen in 1997 (kg N):

 (-) None
 (N) 160

Experimental diary:

22-Sep-97 : B : Ploughed.
25-Sep-97 : B : Rotary harrowed, Genesis, dressed Sibutol, drilled at 380 seeds per m².
26-Sep-97 : B : Rolled.
12-Nov-97 : B : Stefes IPU 500 at 2.0 l with Stomp 400 SC at 2.0 l and Cyperkill 10 at 250 ml in 200 l.
17-Feb-98 : B : 34.5% N at 120 kg.
17-Apr-98 : B : 34.5% N at 460 kg.
08-May-98 : B : Ally at 20 g with Pointer at 0.5 l in 200 l.
28-May-98 : B : Opus at 0.7 l in 200 l.
20-Aug-98 : B : Combine harvested.

98/R/CS/457

NOTE: Plant samples were taken in July to assess root and stem base diseases.

GRAIN TONNES/HECTARE

***** Tables of means *****

CROP	(R)	(W)	Mean
SETDESTR			
(PG)	9.62	5.19	7.40
(PC)	9.59	5.75	7.67
(MP)	7.74	5.52	6.63
(HP)	9.32	5.36	7.34
(-P)	8.95	4.94	6.95
Mean	9.04	5.35	7.20

N	(-)	(N)	Mean
SETDESTR			
(PG)	7.18	7.63	7.40
(PC)	8.08	7.25	7.67
(MP)	6.66	6.59	6.63
(HP)	7.64	7.04	7.34
(-P)	6.81	7.08	6.95
Mean	7.27	7.12	7.20

N	(-)	(N)	Mean
CROP			
(R)	8.95	9.13	9.04
(W)	5.60	5.10	5.35
Mean	7.27	7.12	7.20

SETDESTR	CROP	(R)	(W)	
	N	(-)	(N)	(N)
(PG)		9.54	9.69	4.81
(PC)		9.64	9.54	6.53
(MP)		7.24	8.24	6.09
(HP)		9.57	9.06	5.70
(-P)		8.77	9.14	4.86

98/R/CS/457

GRAIN TONNES/HECTARE

*** Standard errors of differences of means ***

SETDESTR	CROP	N	SETDESTR CROP
0.542	0.343	0.186	0.766
SETDESTR N	CROP N	SETDESTR CROP N	
0.616	0.390	0.872	
Except when comparing means with the same level(s) of			
SETDESTR	0.415		
CROP	0.263		
SETDESTR.CROP		0.587	

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	18	0.939	13.0
BLOCK.WP.SP	20	0.719	10.0

GRAIN MEAN DM% 86.8

SUB-PLOT AREA HARVESTED 0.00230