


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1999

[Full Table of Content](#)



99/R/LNS/1 Fungicides and Spring Linseed - Linseed

Rothamsted Research

Rothamsted Research (2000) *99/R/LNS/1 Fungicides and Spring Linseed - Linseed* ; Yields Of The Field Experiments 1999, pp 149 - 151 - DOI: <https://doi.org/10.23637/ERADOC-1-34>

99/R/LNW/2

CLEAN GRAIN TONNES/HECTARE (WEED SEED REMOVED)

*** Standard errors of differences of means ***

WEEDSPEC	WEEDDENS	TIMWEED	WEEDSPEC WEEDDENS
0.082	0.116	0.082	0.164
WEEDSPEC TIMWEED	WEEDDENS TIMWEED	WEEDSPEC WEEDDENS TIMWEED	
0.116	0.164	0.231	

S.e.d. for comparing NO with any item in the WEEDSPEC.WEEDDENS.TIMWEED table is 0.200

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	35	0.283	12.4

PLOT AREA HARVESTED 0.00230

99/R/LNS/1

LINSEED

FUNGICIDES AND SPRING LINSEED

Object: To test fungicidal spray programmes on spring linseed - Sawyers I.

Sponsors: B.D.L. Fitt, S.A.M. Perryman.

Design: 2 blocks of 2 plots randomly split into 3 x 4 + 2.

Plot dimensions: 3.0 x 12.0.

Treatments: All combinations of:-

Whole plots: All combinations of:-

1. VARIETY

A	Antares
J	Jupiter

Sub-plots

2. FUNGICIDE Target organism and fungicides used:

AL	Alternaria (Rovral Flo at 2.0 l in 220 l)
BO	Botrytis (Benlate Fungicide at 1.1 kg in 220 l)
A+B	Alternaria and Botrytis (Compass at 3.0 l in 220 l)

3. TIMING Fungicide timing:

P+M	Pre-flowering
MF	Mid-flowering
CD	Capsule development
M+C	Mid-flowering and capsule development

Plus two extra treatments:

EXTRA

A	Antares with no fungicide (duplicated)
J	Jupiter with no fungicide (duplicated)

Experimental diary:

03-Dec-98	: B :	: Ploughed, started.
04-Dec-98	: B :	: Ploughed, completed.
11-Feb-99	: B :	: PK as (0:20:32) at 1000 kg.
29-Mar-99	: B :	: Spring-tined.
29-Mar-99	: T : A	: Combination drilled, Antares, tr. Vitavax RS, at 700 seeds/m ² with the Accord drill.
29-Mar-99	: T : J	: Combination drilled, Jupiter, tr. Hydraguard and Prelude 20 LF, at 700 seeds/m ² with the Accord drill.
28-Apr-99	: B :	: 34.5% N at 174 kg.

99/R/LNS/1

Experimental diary:

14-May-99 : B : : Ally at 30 g in 200 l.
 09-Jun-99 : T : P+M : Fungicides applied.
 25-Jun-99 : T : M+C,MF,P+M : Fungicides applied.
 13-Jul-99 : T : CD,M+C : Fungicides applied.
 29-Jul-99 : B : : tm) Enhance at 400 ml in 400 l.
 : B : : tm) Reglone at 3.0 l in 400 l.
 01-Sep-99 : B : : Combine harvested.

Previous crops: S. barley 1997, w. wheat 1998.

GRAIN TONNES/HECTARE

***** Tables of means *****

FUNGICIDE	AL	BO	A+B	Mean
TIMING				
P+M	2.17	2.31	2.28	2.26
MF	2.14	2.16	2.32	2.21
CD	1.98	2.13	2.03	2.05
M+C	2.12	2.06	2.30	2.16
Mean	2.10	2.17	2.24	2.17

VARIETY	A	J	Mean
TIMING			
P+M	2.26	2.26	2.26
MF	2.20	2.22	2.21
CD	1.97	2.12	2.05
M+C	2.15	2.17	2.16
Mean	2.14	2.19	2.17

VARIETY	A	J	Mean
FUNGICIDE			
AL	2.06	2.15	2.10
BO	2.18	2.15	2.17
A+B	2.20	2.27	2.24
Mean	2.14	2.19	2.17

FUNGICIDE	AL		BO		A+B	
VARIETY	A	J	A	J	A	J
TIMING						
P+M	2.15	2.19	2.28	2.35	2.33	2.24
MF	2.06	2.22	2.22	2.11	2.31	2.33
CD	1.92	2.04	2.06	2.20	1.94	2.12
M+C	2.08	2.15	2.17	1.95	2.21	2.40

EXTRA	A	J	Mean
	1.96	2.11	2.03

Grand mean 2.15