

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1999

[Full Table of Content](#)



99/R/CS/476 Fungicide Sequence and Take-all - W. Wheat

Rothamsted Research

Rothamsted Research (2000) *99/R/CS/476 Fungicide Sequence and Take-all - W. Wheat* ; Yields Of The Field Experiments 1999, pp 83 - 84 - DOI: <https://doi.org/10.23637/ERADOC-1-34>

99/R/CS/472

GRAIN TONNES/HECTARE

***** Tables of means *****

CROP FUNGIDE	WW	BW	Mean
---	8.50	7.74	8.12
E-E	8.26	8.00	8.13
-E-	7.56	7.39	7.48
EEE	8.49	7.85	8.17
B-B	8.71	7.76	8.23
-B-	7.54	7.94	7.74
BBB	8.31	8.13	8.22
Mean	8.23	7.82	8.03

*** Standard errors of differences of means ***

FUNGIDE	CROP	FUNGIDE	CROP
0.261		0.369	min.rep
0.226	0.131	0.320	max-min
		0.261	max.rep

FUNGIDE
max.rep - only
max-min - v any of the remainder
min.rep Any of the remainder

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	46	0.522	6.5
GRAIN MEAN DM%	87.6		
PLOT AREA HARVESTED	0.00227		

99/R/CS/476

FUNGICIDE SEQUENCES AND TAKE-ALL

Object: To determine the effects of a fungicidal seed treatment on take-all (*Gaeumannomyces graminis*) development in w. wheat - Long Hoos IV 4.

Sponsors: J.F. Jenkyn, R.G. Gutteridge, G.L. Bateman.

The third year, w. wheat.

For previous years see 97-98/R/CS/476.

Design: 4 randomised blocks of 2 x 2 x 2.

Plot dimensions: 3.0 x 10.0.

Treatments: All combinations of:-

- | | |
|-----------|---|
| 1. FUNG97 | Fungicidal seed treatment to the 1997 crop: |
| (F97) | Seed treated |
| (-97) | None |
| 2. FUNG98 | Fungicidal seed treatment to the 1998 crop: |
| (F98) | Seed treated |
| (-98) | None |
| 3. FUNG99 | Fungicidal seed treatment to the 1999 crop: |
| F99 | Seed treated |
| -99 | None |

NOTE: The seed treatment was fluquinconazole at 75 g/100 kg of seed.

Experimental diary:

26-Sep-98	: B :	: Ploughed.
12-Oct-98	: B :	: Spring-tined.
	: B :	: Combination drilled, Hereward, tr as treatment, at 380 seeds per m ² with the Accord drill.
	: B :	: Hardy at 7.5 kg.
09-Mar-99	: B :	: tm)Isoguard at 1.0 l in 200 l.
	: B :	: tm)Unite A at 0.125 l in 200 l.
	: B :	: tm)Unite B at 1.0 l in 200 l.
	: B :	: tm)Cropoil at 1.0 l in 200 l.
12-Mar-99	: B :	: 34.5% N at 145 kg.
14-Apr-99	: B :	: 34.5% N at 435 kg.
29-Jul-99	: B :	: Combine harvested.