This work is licensed under a Creative Commons Attribution 4.0 International License.

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.

Yields of the Field Experiments 2013

Full Table of Content

R/CS/477 & W/CS/478 Continuous Maize

Rothamsted Research

Rothamsted Research (2014) *R/CS/477 & W/CS/478 Continuous Maize ;* Yields Of The Field Experiments 2013, pp 52 - 57 **- DOI: https://doi.org/10.23637/ERADOC-1-223**

This work is licensed under a Creative Commons Attribution 4.0 International License.

13/R/CS/477

CONTINUOUS MAIZE

Object: To monitor the fate of organic carbon in the soil organic matter – Hoosfield

Sponsors: A. J. Macdonald

The 17th year, forage maize and s. barley

For previous years see Yield Books for 97-12/R/CS/477

Design: 3 randomised blocks of 6 plots.

Plot dimensions: 12.0 x 25.0

Treatments:-

CROP Crop and straw treatments:

M	Continuous maize, stubble incorporated
(M)B	S. barley after five years maize, stubble incorporated
MT	Maize, stubble plus 10 t maize tops incorporated
B(M)	S. barley, after ten years of Maize, straw removed
BT	Continuous spring barley, straw removed plus 10 t maize tops incorporated
B	Continuous spring barley, straw removed

Note: Cropping was changed from Maize to S. barley on the BM treatment in 2010

Experimental diary

Date		Application	Rate	Units
28-Sept-12	f	TSP applied – all plots	171	kg/ha
		MOP applied – all plots	181	kg/ha
05-Oct-12	a	Maize tops spread on plots 3, 9,18, 6, 12, 16	10	t/ha
02-Apr-13	a	Spring tined		
03-Apr-13	s	Drilled Barley only, var. Tipple dr Rancona	350	seeds/m^2
6-Apr-13	p	Sprayed Kula	3.5	l/ha
16-May-13	a	flexitined maize plots	—	—
20-May-13	a	Power-harrowed maize plots	—	—
20-May-13	a	Drilled Maize, Hudson tr Mesurol	as plan	
21-May-13	f	Applied Doubletop to Maize and Barley	@356	kg/ha
26-May-13	p	Sprayed Refine Max, Compitore Plus, Mobius, Cyflomid	re@75 co@1.0 mo@0.6 cyf@0.125	g/ha g/ha l/ha l/ha l/ha
26-Jun-13	p	Sprayed Mobius	@0.4	l/ha
26-Jun-13	p	Sprayed Samson and Callisto, Maize sprayed only	Both @0.5	l/ha
10-Jul-13	a	Pulling Wild Oats	—	—
19-Jul-13	a	Cut Paths.	—	—
12-Aug-13	a	Claas - Harvested opened up exp.	—	—

https://doi.org/10.23637/ERADOC-1-223

This work is licensed under a Creative Commons Attribution 4.0 International License.

27-Aug-13	a	Claas - Harvested OE's	—	—
27-Aug-13	a	Sampo - Harvested all plots	—	—
29-Aug-13	a	Claas - Harvested, cleared OE's	—	—
25-Sep-13	a	Harvested all Maize plots	—	—
25-Sep-13	a	Cleared OE's Maize	—	—
26-Sep-13	a	Cut Maize OE's	—	—

NOTE: Samples of barley grain and maize (whole crop) were taken for chemical analyses.

```
MAIZE

WHOLE CROP TONNES/HECTARE (100% DM)

***** Tables of means *****

    Treatment
            M      2.33
           MT      2.72
         M(B)      2.94
         (B)M      1.87

         Mean      2.47

Standard errors of differences of means
---------------------------------------

Table            Treatment
rep.                     3
d.f.                     6
s.e.d.               0.678

Stratum standard errors and coefficients of variation
=====================================================

Variate: TPlDm Total plant dry matter tonnes/hectare

Stratum                   d.f.          s.e.         cv%

Blocks                       2         0.437        17.7
Blocks.Plots                 6         0.830        33.7

MEAN DM% 23.8

Plot area harvested 0.00108
```

Note: Maize yields were adversely affected by the accidental application of residual herbicide (Topik). Therefore, yields are unreliable.

https://doi.org/10.23637/ERADOC-1-223

This work is licensed under a Creative Commons Attribution 4.0 International License.

13/R/CS/477

SPRING BARLEY

Grain tonnes/hectare

***** Tables of means *****

Treatment	
BT	5.10
B	4.85
Mean	4.98

Standard errors of differences of means

Table	Treatment
rep.	3
d.f.	2
s.e.d.	0.086

Stratum standard errors and coefficients of variation
==========

Variate: Grain85% Grain (at 85% dry matter) tonnes/hectare

Stratum	d.f.	s.e.	cv%
Blocks	2	0.299	6.0
Blocks.Plots	2	0.106	2.1

GRAIN MEAN DM% 87.1

Plot area harvested 0.00525

https://doi.org/10.23637/ERADOC-1-223

This work is licensed under a Creative Commons Attribution 4.0 International License.

13/W/CS/478

CONTINUOUS MAIZE

Object: To monitor the fate of organic carbon in the soil organic matter – Woburn, Stackyard AI

Sponsors: A. J. Macdonald

The 17th year, forage maize and s. barley

For previous years see Yield Books for 97-12/W/CS/478

Design: 3 randomised blocks of 6 plots.

Plot dimensions: 9.0 x 25.00

Treatments:-

CROP Crop and straw treatments:

M Continuous maize, stubble incorporated
(M)B S. barley after five years maize, stubble incorporated
MT Maize, stubble plus 10 t maize tops incorporated
B(M) S. barley, after ten years of maize, straw removed
BT Continuous spring barley, straw removed plus 10 t maize tops incorporated
B Continuous spring barley, straw removed
Note: Cropping was changed from Maize to S. barley on the BM treatment in 2010
NOTE: Samples of barley grain and maize (whole crop) were taken for chemical analyses.

Experimental diary

Date		Application	Rate	Units
23-Oct-12	a	Applied Maize tops, plots 2, 4, 12, 13, 16, 17.	10	t/ha
06-Nov-12	a	Ploughed North	—	—
15-Mar-13	a	Spring tined	—	—
04-Apr-13	s	Drilled NFC Tipple, tr Rancona	350	seeds/m^2
04-Apr-13	a	Rolled, rolled Oats/sp Barley	—	—
20-Apr-13	f	Applied TSP	171	kg/ha
22-Apr-13	f	Applied MOP	181	kg/ha
23-Apr-13	f	Applied Double Top, applied to Sp. Barley and maize seedbed	356	kg/ha
22-May-13	s	Drilled Maize, Hudson tr Mesurol	10.1	seeds/m^2
02-Jul-13	p	Sprayed Harmony M, Mobius and Hatchet Xtra, sprayed spring barley only	Har @ 0.1 Mob @ 0.5 Hat @ 0.7	kg/ha l/ha l/ha
05-Jul-13	p	Sprayed Mobius, sprayed spring barley only	0.4	l/ha
05-Jul-13	p	Sprayed Samson and Callisto, Maize plots only	0.5 0.5	l/ha l/ha
31-Aug-13	a	Cut plots for yield	—	—

https://doi.org/10.23637/ERADOC-1-223

This work is licensed under a Creative Commons Attribution 4.0 International License.

04-Sep-13	a	Combined	—	—
06-Sep-13	a	Baled	—	—
07-Oct-13	a	Cut Maize for yields	—	—
07-Oct-13	a	Mowed and Baled, only two bales of maize	—	—

```
MAIZE WHOLE CROP TONNES/HECTARE (100% DM)

***** Tables of means *****

    Treatment
         M      2.79
         T      4.22
      M(B)      3.11
      (B)M      3.34

       Mean     3.36

Standard errors of differences of means
---------------------------------------

Table            Treatment
rep.                     3
d.f.                     6
s.e.d.               0.628

Stratum standard errors and coefficients of variation
=====================================================
Stratum                  d.f.          s.e.         cv%

Blocks                      2         0.390        11.6
Blocks.Plots                6         0.769        22.8

Mean DM% 26.5

Plot area harvested 0.00108
```


https://doi.org/10.23637/ERADOC-1-223

This work is licensed under a Creative Commons Attribution 4.0 International License.

13/W/CS/478

```
SPRING BARLEY

GRAIN TONNES/HECTARE

***** Tables of means *****

    Treatment
           BT      3.93
            B      3.06

         Mean      3.49

Standard errors of differences of means
---------------------------------------

Table            Treatment
rep.                     3
d.f.                     2
s.e.d.               0.247

Stratum standard errors and coefficients of variation
=====================================================

Stratum                  d.f.          s.e.         cv%

Blocks                      2         0.203         5.8
Blocks.Plots                2         0.303         8.7

Grain mean DM%  88.0

Plot area harvested  0.00525

Standard errors of differences of means
---------------------------------------
Table            Treatment
s.e.d.               0.095

Stratum standard errors and coefficients of variation
=====================================================
Stratum                  d.f.          s.e.         cv%

Blocks.Plots                4         0.116         2.4

GRAIN MEAN DM% 84.3

PLOT AREA HARVESTED 0.00525
```


https://doi.org/10.23637/ERADOC-1-223