


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1957

[Full Table of Content](#)



57/R/CH/1 Lucerne - Single and Repeated Applications of K - 3rd Year

Rothamsted Research

Rothamsted Research (1958) *57/R/CH/1 Lucerne - Single and Repeated Applications of K - 3rd Year ; Yields Of The Field Experiments 1957*, pp 114 - 115 - DOI:

<https://doi.org/10.23637/ERADOC-1-177>

57/Ch/1.1

LUCERNE

Single and repeated applications of potash - Great Harpenden II 1957,
the third year.

Design: 6 randomized blocks of 8 plots each.

Area of each plot: 0.0147 acres.

Treatments: (For the 3 years 1955 - 57).

None (2 plots per block).

Single dressings in 1955: 1; 2; 3 cwt K_2O per acre as muriate
of potash.

Annual dressings: 0.33; 0.66; 1.0 cwt K_2O per acre as muriate
of potash.

Note. In spring 1955 all the potash dressings were applied in the
seedbed. Subsequent annual applications are given as top dressings.

Basal dressing: None.

Cultivations, etc.: Annual potash dressings applied: Dec 3, 1956.

Cut and weighed green: three times - May 6, 1957, July 8, Oct 15.

Standard errors per plot. Dry matter:

1st cut:	1.87 cwt per acre or 7.6% (36 d.f.)
2nd cut:	2.40 cwt per acre or 6.0% (36 d.f.)
3rd cut:	1.33 cwt per acre or 8.0% (36 d.f.)
Total of 3 cuts:	3.44 cwt per acre or 4.2% (36 d.f.)

Note. For first and second year's results, see "Results of the Field
Experiments" 1955, Section Cc/1; 1956, Section Cg/1.

57/Ch/1.2

Summary of Results

Dry matter: cwt per acre

		K ₂ O: cwt per acre							
		Applied 1955			Applied annually 1955, 1956 & 1957				
	None	1.0	2.0	3.0	0.33	0.66	1.0	Mean	
1st cut									
	(±0.54)	(±0.76)							
Mean	22.9	23.3	25.2	26.1	24.6	25.6	27.0	24.7	
Increase (±0.93)		+0.4	+2.3	+3.2	+1.7	+2.7	+4.1		
2nd cut									
	(±0.69)	(±0.98)							
Mean	38.3	39.0	41.6	42.3	39.8	41.7	41.4	40.3	
Increase (±1.20)		+0.7	+3.3	+4.0	+1.5	+3.4	+3.1		
3rd cut									
	(±0.39)	(±0.54)							
Mean	16.3	16.8	16.7	17.0	17.1	17.1	16.2	16.7	
Increase (±0.67)		+0.5	+0.4	+0.7	+0.8	+0.8	-0.1		
Total of 3 cuts									
	(±0.99)	(±1.40)							
Mean	77.5	79.1	83.5	85.4	81.5	84.4	84.6	81.7	
Increase (±1.72)		+1.6	+6.0	+7.9	+4.0	+6.9	+7.1		

Mean dry matter % as cut: 1st 25.4
 2nd 32.1
 3rd 33.3
 Total of 3 cuts 30.3