


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1957

[Full Table of Content](#)



57/R/BA/3 and 57/W/BA/3 Six-course Rotation

Rothamsted Research

Rothamsted Research (1958) *57/R/BA/3 and 57/W/BA/3 Six-course Rotation* ; Yields Of The Field Experiments 1957, pp 24 - 27 - DOI: <https://doi.org/10.23637/ERADOC-1-177>

57/Ba/3.1

SIX COURSE ROTATION EXPERIMENT

The 28th year

Seasonal effects of fertilizers - Rothamsted Long Hoos IV and Woburn Stackyard 1957.

For history, treatments, etc., see "Details of the Classical and Long Term Experiments" 1956.

Area of each plot: Rothamsted, 0.0250 acres; Woburn, 0.0266 acres.

Cultivations, etc.:

Rothamsted

Sugar beet.

Ploughed twice: Sept 21, 1956 and Nov 12. Fertilizers applied: Apr 13, 1957. Seed drilled at 12 lb per acre: Apr 16. Sprayed with miscible DDT, 3 pints in 20 gallons per acre: May 29. Singled: June 14 - 19. Sprayed with methyl demeton, 12 oz in 80 gallons per acre: July 9. Lifted: Nov 4 - 13. Variety: Klein E.

Barley.

Sugar beet tops spread: Dec 6, 1956. Ploughed: Jan 12, 1957. Ground chalk applied at 20 cwt per acre: Feb 21. Fertilizers applied: Mar 16. Seed drilled at $2\frac{3}{4}$ bushels per acre: Mar 18. Clover seed undersown: May 20. Harvested: Aug 8. Variety: Plumage Archer.

Clover.

Seed undersown in barley at 40 lb per acre: Apr 23, 1956. Autumn fertilizers applied: Nov 16. Sulphate of ammonia applied: Mar 30, 1957. Cut: July 9. Variety: S123 Late Flowering Red.

Wheat.

Ploughed twice: July 27, 1956 and Sept 21. Seed drilled at $2\frac{3}{4}$ bushels per acre: Oct 25. Autumn fertilizers applied: Oct 26. Sulphate of ammonia applied: Apr 24, 1957. Harvested: Aug 8. Variety: Yeoman.

Potatoes.

Ploughed 3 times: Sept 21, 1956, Nov 12, Jan 22, 1957. Ridged, fertilizers applied: May 3. Potatoes planted: May 4. Earthed up: July 6. Sprayed with copper fungicide, 5 lb in 40 gallons per acre: July 31 and Aug 21. Sprayed with sulphuric acid, 15% BOV at 100 gallons per acre: Sept 19. Lifted: Sept 30. Variety: Majestic.

Rye.

Ground chalk applied at 20 cwt per acre, ploughed: Oct 15, 1956. Seed drilled at 3 bushels per acre: Oct 25. Autumn fertilizers applied: Oct 26. Sulphate of ammonia applied: Apr 24, 1957. Harvested: Aug 7. Variety: King II.

57/Ba/3.2

Woburn

Sugar beet.

Ploughed twice: Sept 1, 1956 and Nov 14. Fertilizers applied: Apr 11, 1957. Seed drilled at 12 lb per acre: Apr 12. Sprayed with miscible DDT, 3 pints in 20 gallons per acre: May 22. Singled: June 12 - 13. Sprayed with methyl demeton, 12 oz in 60 gallons per acre: July 8. Lifted: Nov 12. Variety: Klein E.

Barley.

Ploughed: Nov 14, 1956. Fertilizers applied: Mar 12, 1957. Seed drilled at $2\frac{1}{2}$ bushels per acre: Mar 16. Harvested: Aug 2. Variety: Herta.

Clover.

Ploughed twice: Aug 24, 1956 and Nov 13. Fertilizers applied: Mar 23, 1957. Seed broadcast at 40 lb per acre: Mar 26. Cut: July 5. Variety: Crimsen clover.

Wheat.

Ploughed twice: July 24, 1956 and Sept 29. Autumn fertilizers applied: Oct 15. Seed drilled at $2\frac{1}{2}$ bushels per acre: Oct 20. Sprayed with MCPP at 7 pints in 40 gallons per acre: Apr 6, 1957. Sulphate of ammonia applied: May 1. Harvested: Aug 7. Variety: Yeoman.

Potatoes.

Ploughed twice: Sept 4, 1956 and Nov 13. Ridged, fertilizers applied and potatoes planted: Apr 26, 1957. Sprayed with miscible DDT, 3 pints in 20 gallons: May 22. Sprayed with copper fungicide, 5 lb in 40 gallons per acre: Aug 3, Aug 22 and Sept 4. Sprayed with arsenious compound, 1 gallon in 40 gallons per acre: Sept 16. Lifted: Oct 1. Variety: Majestic.

Rye.

Ploughed: Oct 6, 1956. Ground chalk applied at 20 cwt per acre: Oct 11. Seed drilled at $2\frac{1}{2}$ bushels per acre: Oct 20. Sprayed with MCPP at 7 pints in 40 gallons per acre: Apr 6, 1957. Sulphate of ammonia applied: Apr 18. Harvested: Aug 7. Variety: King II.

Note: In 1957 at Woburn, the nitrogen levels were doubled on all crops except clover.

57/Ba/3.3

Summary of ResultsMean yields per acre and responses in yield per cwt of N, P₂O₅ and K₂O

	Rothamsted	Woburn	Rothamsted	Woburn
Sugar Beet, roots (washed): tons per acre			Barley, grain: * cwt per acre	
Mean	7.71	14.15	26.7	28.4
Response to: N	+2.55	+2.27	+12.9	+9.9
P	-0.80	+0.48	-12.4	+13.6
K	-0.31	+2.75	-4.9	-1.0
Mean dry matter % as harvested:			83.3	**
Sugar Beet, sugar percentage			Barley, straw: * cwt per acre	
Mean	16.7	17.5	16.1	24.3
Response to: N	+0.7	-0.3	+16.9	+10.6
P	+1.1	0.0	-7.2	+7.5
K	+0.8	+1.1	+7.3	+1.5
Mean dry matter % as harvested:			83.1	**
Sugar Beet, total sugar: cwt per acre			Clover, hay, dry matter: cwt per acre	
Mean	25.7	49.8	25.1	8.9
Response to: N	+9.5	+7.1	-14.0	+14.8
P	-0.8	+1.7	+8.7	-5.7
K	+0.1	+12.9	+4.0	-2.1
Mean dry matter % as cut:			58.7	25.3
Sugar Beet, tops: tons per acre			Wheat, grain: * cwt per acre	
Mean	9.26	10.09	32.1	10.0
Response to: N	+5.43	+6.35	+9.4	+2.5
P	+0.69	-0.19	+0.9	-17.6
K	-1.08	+0.13	-4.2	+9.3
Mean dry matter % as harvested:			83.7	**
Sugar Beet, plant number: thousands per acre			Wheat, straw: * cwt per acre	
Mean	29.6	**	43.5	11.8
Response to: N	-1.3		+21.3	+2.2
P	-0.3		-4.4	-17.3
K	+0.3		-8.2	+4.0
Mean dry matter % as harvested:			84.9	**

*(At 85% dry matter). ** Not recorded.

57/Ba/3.4

Mean yields per acre and responses in yield per cwt of N, P₂O₅ and K₂O

	Rothamsted	Woburn	Rothamsted	Woburn
Potatoes, total tubers: tons per acre			Rye, grain: cwt per acre	
Mean	7.38	12.16	*	
Response to: N	+0.15	+2.57	29.2	24.8
P	+1.15	-0.16	+14.9	+14.7
K	+0.61	+0.17	-2.3	+13.5
Mean dry matter % as harvested:			-3.5	-0.9
			82.0	**
Potatoes, percentage ware			Rye, straw: cwt per acre	
	(1)	(2)	*	
Mean	80.0	90.5	32.9	28.9
Response to: N	-4.4	+3.1	+7.2	+10.6
P	+8.3	+2.4	-3.1	+15.5
K	+5.6	-3.0	-5.8	-1.4
Mean dry matter % as harvested:			84.7	**

*(At 85% dry matter).

Riddle: (1) 1½"; (2) 1⅝"