


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1957

[Full Table of Content](#)



57/R/CA/2 Winter Wheat - Varieties and N

Rothamsted Research

Rothamsted Research (1958) *57/R/CA/2 Winter Wheat - Varieties and N* ; Yields Of The Field Experiments 1957, pp 70 - 70 - DOI: <https://doi.org/10.23637/ERADOC-1-177>

57/Ca/2

WINTER WHEAT

Varieties and levels of nitrogen - Great Knott I 1957.

Design: 3 randomized blocks of 8 plots each, plots being split into 2 for the application of nitrogen.

Area of each sub plot: 0.0101 acres. Area harvested: 0.0067 acres.

Treatments: All combinations of:-

Whole plots. Varieties: Banco (1); Cappelle (2); Heine 7 (3); Hybrid 46 (4); Leda (5); Minister (6); Squareheads' Master 13/4 (7); Yeoman (8).

Sub plots. Nitrogen: 0.4; 0.8 cwt N per acre applied as sulphate of ammonia.

Basal dressing: $1\frac{1}{4}$ cwt compound fertilizer (16% P_2O_5 , 16% K_2O) per acre combine drilled with seed.

Cultivations, etc.: Ploughed: Nov 15, 1956. Seed combine drilled at $2\frac{3}{4}$ bushels per acre: Nov 27. Sulphate of ammonia applied: May 1, 1957. Sprayed with DNOC at 8 lb in 80 gallons per acre: May 20. Combine harvested: Aug 25. Previous crop: Spring beans.

Standard errors per plot, Grain (at 85% dry matter):

Whole plot: 2.28 cwt per acre or 6.3% (14 d.f.)

Sub plot: 1.93 cwt per acre or 5.4% (16 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

N: cwt per acre	Variety								Mean	
	1	2	3	4	5	6	7	8		
	(±1.53)*									
0.4	30.5	40.3	34.3	38.2	42.3	36.7	29.4	22.7	34.3	
0.8	33.7	42.4	40.5	42.4	44.1	38.5	33.8	26.6	37.7	
Mean	(±1.32)	32.1	41.3	37.4	40.3	43.2	37.6	31.6	24.7	36.0
Difference	(±1.58)	3.2	2.1	6.2	4.2	1.8	1.8	4.4	3.9	3.4
										(±0.56)

*for use in comparisons other than vertical.

Mean dry matter % as harvested: 83.8