


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1957

[Full Table of Content](#)



57/R/A/4 Barnfield - Mangolds and Sugar Beet

Rothamsted Research

Rothamsted Research (1958) *57/R/A/4 Barnfield - Mangolds and Sugar Beet* ; Yields Of The Field Experiments 1957, pp 8 - 10 - DOI: <https://doi.org/10.23637/ERADOC-1-177>

57/A/4.1

MANGOLDS AND SUGAR BEET - BARNFIELD

The 82nd and 12th years

For history, treatments etc., see "Details of the Classical and Long Term Experiments" 1956.

A maintenance chalk dressing was applied to Series A, AC and C, at 6 cwt, 9 cwt and 3 cwt per acre respectively, under the new liming scheme introduced in spring 1956.

In July, 4 rows of mangolds and 4 of sugar beet on each plot were thinned, and the remainder removed by thistle bar. Plot 4 N(a) was left with 4 rows of mangolds only.

Cultivations, etc.: Ground chalk applied: Nov 30, 1956. Dung applied: Nov 28 - Dec 1. Ploughed: Dec 1 - 5. Fertilizers applied: May 10 - 14, 1957. Sugar beet drilled at 16 - 20 lb per acre; mangolds drilled at 8 lb per acre: May 17. Sprayed the uncropped west side of plots 1N and 10 only before emergence with 2 gallons Pentachlorophenol(PCP) in 40 gallons of water per acre: May 30. Singled: July 9 - 16. Removed discarded rows by thistle bar: July 10. Top dressings applied to reduced plot areas: July 16 - 18. Lifted: Oct 26 - Nov 7. Varieties: Mangolds - Yellow Globe, sugar beet - Klein E.

Cultivations to discarded areas after July 10:-

Thistle bar: July 11, July 25, July 29, Sept 17.

57/A/4.2

Summary of Results

Strip	Cross Dressing				
	O	N	A	AC	C
Mangolds, roots: tons per acre					
1	20.35	31.29	31.46	29.41	25.65
2	20.52	30.78	29.07	33.17	28.04
4	7.02	(a) 18.47*	22.57	27.70	21.37
		(b) 24.45*			
5	5.73	19.32	10.77	13.17	13.34
6	6.03	16.41	13.85	23.77	14.02
7	7.54	16.24	21.54	26.33	26.67
8	3.38	13.34	12.65	13.51	13.17
9	21.88				
Mangolds, leaves: tons per acre					
1	5.45	7.84	8.84	9.67	8.45
2	6.03	8.30	7.89	9.53	7.89
4	2.12	(a) 6.62*	9.18	11.24	9.11
		(b) 7.82*			
5	2.00	5.64	6.45	7.52	6.55
6	2.17	4.25	4.69	8.67	5.69
7	1.98	4.93	6.42	8.99	8.11
8	1.47	4.79	5.18	4.91	4.86
9	7.01				
Mangolds, plant number: thousands per acre					
1	16.1	19.3	18.5	17.8	17.0
2	17.0	19.2	18.1	18.8	17.8
4	18.0	(a) 17.4*	18.5	15.5	15.5
		(b) 18.9*			
5	17.0	19.9	19.1	19.3	19.5
6	18.5	16.6	12.8	15.8	13.2
7	19.1	19.9	18.3	18.6	20.2
8	16.1	20.2	18.2	14.0	16.8
9	20.4				

*No nitrate of soda. Nitrogen applied as calcium and potassium nitrates.

57/4/4.3

Strip	Cross Dressing				
	0	N	A	AC	C
Sugar beet, roots (washed): tons per acre					
1	8.13	11.10	11.39	11.87	12.10
2	7.56	10.64*	11.64	13.73	11.11
4	3.96	(b) 9.35*	10.20	11.42	10.42
5	3.16	8.48	7.27	7.43	7.95
6	3.87	7.28	8.92	10.03	7.79
7	3.40	6.50	9.62	9.49	8.81
8	2.78	6.80	6.72	7.97	6.94
9	7.94				

Sugar beet, tops: tons per acre					
1	7.89	10.89	13.36	13.97	13.80
2	7.77	15.12*	12.80	15.17	13.60
4	3.27	(b) 10.09*	9.38	15.09	13.63
5	2.56	10.06	10.11	12.48	11.55
6	2.42	7.99	9.13	12.87	9.13
7	2.69	8.89	9.70	14.70	11.92
8	2.39	9.45	8.96	13.46	10.01
9	8.23				

Sugar beet, plant number: thousands per acre					
1	18.1	19.0	19.6	20.1	18.8
2	19.2	20.4*	20.8	19.1	18.7
4	24.5	(b) 21.0*	21.7	18.0	18.9
5	23.6	23.0	21.3	19.6	19.8
6	25.1	21.3	21.9	18.2	18.2
7	24.9	22.4	22.3	19.4	20.5
8	22.5	22.7	22.4	20.0	19.7
9	23.1				

Sugar beet, sugar percentage					
1	17.9	16.6	16.7	15.5	16.2
2	17.7	16.0*	16.5	15.7	17.0
4	18.5	(b) 16.8*	17.2	15.3	16.7
5	18.1	16.4	17.1	15.9	15.7
6	18.2	16.8	17.2	16.3	16.4
7	18.9	17.1	17.0	15.0	17.0
8	18.4	16.7	17.2	15.9	16.4
9	17.4				

* No nitrate of soda. Nitrogen applied as calcium and potassium nitrates.