


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1957

[Full Table of Content](#)



57/R/CC/1 Spring Oats - Varieties and N

Rothamsted Research

Rothamsted Research (1958) *57/R/CC/1 Spring Oats - Varieties and N* ; Yields Of The Field Experiments 1957, pp 87 - 87 - DOI: <https://doi.org/10.23637/ERADOC-1-177>

57/Cc/1

SPRING OATS

Varieties and levels of nitrogen - Fosters 1957.

Design: 3 randomized blocks of 8 plots each, plots being split into 2 for the application of nitrogen.

Area of each sub plot: 0.0212 acres. Area harvested: 0.0141 acres.

Treatments: All combinations of:-

Whole plots. Varieties: Blenda (1); Deva (2); Flamande (3);
Opus (4); Palu (5); Pendek (6); Sun II (7); de Wattines (8).
Sub plots. Nitrogen: None; 0.36 cwt N per acre applied as
'Nitro-Chalk'.

Basal dressing: 3 cwt compound fertilizer (12% N, 9% P_2O_5 , 9% K_2O) per acre combine drilled with seed.

Cultivations, etc.: Ploughed: Nov 6, 1956 and again Jan 19, 1957.
'Nitro-Chalk' applied: Mar 22. Seed combine drilled at $3\frac{1}{2}$ bushels per acre: Mar 23. Sprayed with DNOC at 8 lb in 80 gallons per acre: May 21. Combine harvested: Aug 8. Previous crop: Wheat.

Standard errors per plot, Grain (at 85% dry matter):

Whole plot: 0.85 cwt per acre or 2.8% (14 d.f.)
Sub plot: 0.80 cwt per acre or 2.6% (16 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

N: cwt per acre (including basal)	1	2	3	Variety		6	7	8	Mean
				4	5				
	(± 0.58)*								
0.36	30.5	29.9	29.0	30.1	30.2	30.9	30.9	28.3	30.0
0.72	32.6	32.1	31.0	32.3	32.0	30.6	31.0	29.6	31.4
Mean (± 0.48)	31.6	31.0	30.0	31.2	31.1	30.7	30.9	29.0	30.6
Difference (± 0.65)	+2.1	+2.2	+2.0	+2.2	+1.8	-0.3	+0.1	+1.3	+1.4 (± 0.23)

*for use in comparisons other than vertical.

Mean dry matter % as harvested: 83.9