


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1957

[Full Table of Content](#)



57/R/CD/1 Cereals and Beans - Rotations

Rothamsted Research

Rothamsted Research (1958) *57/R/CD/1 Cereals and Beans - Rotations* ; Yields Of The Field Experiments 1957, pp 88 - 88 - DOI: <https://doi.org/10.23637/ERADOC-1-177>

57/Ca/1

CEREALS AND BEANS ROTATIONS

The effect of crop sequences on the incidence of cereal foot and root rot diseases - Great Field I 1957 - the 1st year.

Design: Three series each of 3 randomized blocks of 6 plots, starting in each of the years 1957, 1958 and 1959. In 1957 there were 3 plots per block with winter wheat.

Area of each plot: 0.0305 acres. Area harvested: 0.0201 acres.

Treatments:

Crop sequences for each series:

1st year:	WW	WW	WW	SW	O	B
2nd year:	WW	O	O	WW	WW	WW
3rd year:	SW	SW	Be	SW	SW	B

WW = Winter wheat, SW = Spring wheat, O = Oats, B = Barley, Be = Beans.

In the 4th year plots will be split for N, and all cropped with winter wheat.

Basal dressing: 23 cwt ground chalk per acre, 2 cwt compound fertilizer (16% P_2O_5 , 16% K_2O) per acre combine drilled with seed.

Nitrogen for cereals: 3 cwt 'Nitro-Chalk' per acre to spring wheat and 2 cwt 'Nitro-Chalk' per acre to oats and barley all in seedbed. 6 cwt 'Nitro-Chalk' per acre to winter wheat as spring top dressing, half applied in March and half in May.

Cultivations, etc.: Ground chalk applied: Nov 6, 1956. Ploughed: Nov 12. Winter wheat combine drilled at $2\frac{1}{2}$ bushels per acre: Nov 15. 1st application of 'Nitro-Chalk' to winter wheat, 'Nitro-Chalk' applied for oats and seed combine drilled at 4 bushels per acre: Mar 13, 1957. 'Nitro-Chalk' applied for barley and seed combine drilled at 2 bushels per acre: Mar 14. 'Nitro-Chalk' applied for spring wheat and seed combine drilled at 3 bushels per acre: Mar 15. 2nd application of 'Nitro-Chalk' to winter wheat: May 2. Sprayed with DNOC at 8 lb in 80 gallons per acre: May 10. Combine harvested: Aug 26. Varieties: Winter wheat - Heine7; spring wheat - Koga II; oats - Sun II; barley - Proctor. Previous crop: Spring beans.

Summary of Results

Yields (at 85% dry matter): cwt per acre

	WW	SW	B	O
Grain	45.5	39.7	40.7	27.3
Straw	26.1	25.3	20.1	22.5

Mean dry matter % as harvested: Grain, 83.5; Straw, 87.2