


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1957

[Full Table of Content](#)



57/R/CF/2 Potatoes - Control of Blight (Copper Spray)

Rothamsted Research

Rothamsted Research (1958) *57/R/CF/2 Potatoes - Control of Blight (Copper Spray)* ; Yields Of The Field Experiments 1957, pp 96 - 96 - DOI: <https://doi.org/10.23637/ERADOC-1-177>

57/Cf/2.1

POTATOES

The control of blight (Phytophthora infestans) by copper fungicide spray - Highfield IV 1957.

Design: 8 randomized blocks of 2 plots each.

Area of each plot: 0.1273 acres. Area harvested: Whole plots - 0.0141 acres; sub plots - 0.0071 acres.

Treatments.

Whole plots: No spray; copper fungicide sprayed twice.

Sub plots: On sprayed plots only, 2 rows damaged by 2 passages of the tractor during copper spray operations were compared with 2 undamaged rows.

Basal dressing: 15 cwt compound fertilizer (9% N, 9% P₂O₅, 15% K₂O) per acre.

Cultivations, etc.: Ploughed twice: Nov 22, 1956 and Feb 22 - Mar 1, 1957. Basal fertilizer applied on flat: Apr 24. Potatoes machine planted: Apr 25 - 26. Earthed up: July 9. Fungicide treatment, 5 lb in 40 gallons per acre, applied twice: July 31 and Aug 21. Sprayed with sulphuric acid, 20% BOV, 100 gallons per acre: Oct 3. Lifted: Oct 14 - 15. Variety: Majestic. Previous crop: Wheat.

Standard errors per plot. Total tubers:

Whole plots: 1.72 tons per acre or 11.8% (7 d.f.)

Sub plots (sprayed plots only): 1.14 tons per acre or 7.6% (7 d.f.)

Note: Estimates were made of the rate of bulking, destruction of foliage by blight, amount of blight on the tubers. 2 rows of King Edward were planted on each plot for an assessment of blight infected tubers.