


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/W/CB/3 Barley - Forms and Application of N

Rothamsted Research

Rothamsted Research (1961) *60/W/CB/3 Barley - Forms and Application of N* ; Yields Of The Field Experiments 1960, pp 100 - 101 - DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/Cb/3.1

BARLEY

Forms of N and methods of application - Butt Close Woburn 1960.

Design: $4 \times 2 \times 4$ in 6 blocks of 16, with certain high order interactions partially confounded with block differences, plus 2 control plots per block.

Area of each plot: 0.0146 acres.

Treatments: No nitrogen and all combinations of:-

<u>Forms of N:</u>	Ammonium sulphate 21% N	(S)
	Calcium nitrate 15.5% N	(C)
	Ammonium nitrate 23% N	(A)
	or Urea 45.6% N	(U)

Levels of N: 0.3; 0.6 cwt N per acre.

Methods of application: Broadcast (B), combine drilled (D), side band placed (P), top dressed (T).

Basal dressing: 2 cwt granular compound fertiliser (14% P_2O_5 , 28% K_2O) per acre cross drilled.

Cultivations, etc.: Ploughed Jan 27 - Feb 8, 1960. Seed drilled at 2 bushels per acre and seedbed fertilisers applied: Mar 24.

Nitrogen top dressings applied: Apr 29. Sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: May 9. Combine harvested: Aug 23. Variety: Proctor. Previous crop: Sugar beet.

Standard error per plot.

Grain (at 85% dry matter): 1.95 cwt per acre or 7.7% (67 d.f.)

60/Cb/3.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

	Form of N				Mean
	S	C	A	U	
Mean (± 0.40)	26.2	26.1	26.2	26.0	26.1 (± 0.20)
N: cwt per acre					
0.3 (± 0.56)	23.3	24.4	24.0	24.4	24.0
0.6	29.1	27.8	28.5	27.7	28.3
Diff. (± 0.80)	+5.8	+3.4	+4.5	+3.3	+4.3 (± 0.40)
Method of application					
B	25.4	24.3	25.6	25.4	25.2
D (± 0.80)	26.3	26.0	25.3	25.5	25.7 (± 0.40)
P	24.5	25.8	24.6	24.9	24.9
T	28.4	28.3	29.6	28.4	28.7

	Method of application			
	B	D	P	T
N: cwt per acre				
0.3 (± 0.56)	22.8	23.8	23.1	26.3
0.6	27.5	27.7	26.8	31.0
Diff. (± 0.80)	+4.7	+3.9	+3.7	+4.7

Control: 18.0 (± 0.56)

General mean: 25.2

Mean dry matter % as harvested (all plots): 81.4

Form of N

S = Ammonium sulphate 21% N

C = Calcium nitrate 15.5% N

A = Ammonium nitrate 23% N

U = Urea 45.6% N

Method of application

B = Broadcast

D = Combine drilled

P = Side band placed

T = Top dressed