


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/R/CD//2 and 60/W/CD/2 Wheat, Barley and Multiple Crops - Residuals of Weedkillers

Rothamsted Research

Rothamsted Research (1961) *60/R/CD//2 and 60/W/CD/2 Wheat, Barley and Multiple Crops - Residuals of Weedkillers* ; Yields Of The Field Experiments 1960, pp 108 - 112 - **DOI:**
<https://doi.org/10.23637/ERADOC-1-180>

60/Cd/2.1

WHEAT, BARLEY AND MULTIPLE CROPS

Residual effects of triazine weedkillers - Rothamsted (R) Great Field I and Great Knott I and Woburn (W) Broad Mead I and Great Hill 1960.

Design: Strip cropping on sites of 1959 experiments:-

Great Field I (R), Great Knott I (R) and Broad Mead I (W): Winter wheat, kale, sugar beet, barley and oats.

Great Hill (W): Spring wheat, kale, sugar beet, barley and oats.

Area of each plot (acres):

Great Field I (R) and Great Knott I (R) - winter wheat; Broad Mead I (W) - winter wheat: 0.0318. Area harvested: 0.0152. All other crops on above fields: 0.0079. Area harvested: 0.0035 - 0.0053. Great Hill (W) - Barley: 0.0393. Area harvested: 0.0170. Other crops on Great Hill: 0.0098. Area harvested: Oats - 0.0043, Sugar beet - 0.0051.

Treatments: Applied in 1959. See 'Results of the Field Experiments' 1959 pages 59/Ce/2 and 59/Cf/5.

Basal dressings per acre:

Oats, barley and spring wheat (all fields): 3 cwt compound fertiliser (16% N, 9% P_2O_5 , 9% K_2O) combine drilled.

Kale and sugar beet (all fields): 10 cwt compound fertiliser (10% N, 10% P_2O_5 , 18% K_2O).

Winter wheat:- Great Field I (R): $1\frac{1}{2}$ cwt compound fertiliser (6% N, 15% P_2O_5 , 15% K_2O) combine drilled and 4 cwt sulphate of ammonia top dressed. Great Knott I (R): $2\frac{1}{2}$ cwt compound fertiliser (6% N, 15% P_2O_5 , 15% K_2O) combine drilled and 5 cwt sulphate of ammonia top dressed. Broad Mead I (W): $2\frac{1}{2}$ cwt compound fertiliser (6% N, 15% P_2O_5 , 15% K_2O) combine drilled and 3 cwt 'Nitro-Chalk' 21 top dressed.

Cultivations, etc.:

Rothamsted. Great Field I (F) and Great Knott I (K). Ploughed: (K) - Oct 9, 1959, (F) - Oct 21. Winter wheat combine drilled at $2\frac{3}{4}$ bushels per acre: (K) - Oct 23, (F) - Oct 26. Barley combine drilled at 2 bushels per acre: Mar 7, 1960. Basal fertiliser applied for kale and sugar beet: Mar 24. Sugar beet drilled at 19 lb per acre: (K) - Apr 6, (F) - Apr 7. Kale drilled at 3 lb per acre: (K) - Apr 8, (F) - Apr 9. Top dressing of sulphate of ammonia applied to winter wheat: (K) - Apr 8, (F) - Apr 12. Winter wheat sprayed with CMPP at 6 pints in 40 gallons per acre: (F) - Apr 21, (K) - Apr 22. Barley and oats sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: (F) - May 6, (K) - May 10. Sugar beet singled: (F) - May 23, (K) - May 25. Sugar beet sprayed with demeton methyl at 12 fluid oz in 60 gallons per acre: May 30. Barley and oats combine harvested: Aug 16. Winter wheat combine harvested: (F) - Aug 23, (K) - Aug 28. Sugar beet lifted: Oct 25. Kale harvested: (F) - Oct 25, (K) - Nov 24.

60/cd/2.2

Woburn. Broad Mead I (B) and Great Hill (G): Ploughed: (B) - Nov 2, 1959. Winter wheat combine drilled at 3 bushels per acre: Nov 11. Ploughed: (G) - Jan 5, 1960. Seed combine drilled: Barley at $2\frac{1}{4}$ bushels, oats at 4 bushels per acre: (B) - Mar 19, (G) - Mar 26; spring wheat at $2\frac{3}{4}$ bushels per acre: (G) - Mar 26. 'Nitro-Chalk' applied to winter wheat: Apr 5. Basal fertiliser applied to kale and sugar beet: (B) - Apr 11, (G) - Apr 14. Kale and sugar beet seed drilled: Apr 14. Kale and sugar beet sprayed with miscible DDT (against flea beetle) at 3 pints in 40 gallons per acre: May 6. Sugar beet singled: May 30. Sugar beet sprayed with demeton methyl at 12 fluid oz in 40 gallons per acre: June 1. Spring wheat, barley and oats combine harvested: (G) - Aug 22. Winter wheat, barley and oats combine harvested: (B) - Sept 8. Sugar beet lifted: Oct 5. Kale harvested: Nov 2.

Varieties (all fields): Winter wheat: Cappelle; spring wheat: Jufy I; barley: Proctor; oats: Condor; sugar beet: Klein E; kale: Thousand head.

Previous crops: Great Field I (R) and Great Hill (W): Potatoes, Great Knott I (R) and Broad Mead I (W): Spring beans.

Note: Owing to damage by birds, no yields were taken for kale and spring wheat on Great Hill (W), nor for barley and oats on Broad Mead I (W).

Summary of Results

Great Field I (R)

	None	S1	Spray in 1959		S3	A2	Mean
			S2				
	<u>Wheat, grain (at 85% dry matter): cwt per acre</u>						
Mean	53.1	51.4	52.5		49.7	52.8	51.9
	<u>Barley, grain (at 85% dry matter): cwt per acre</u>						
Mean	46.1	44.2	47.5		42.3	44.5	45.1
	<u>Oats, grain (at 85% dry matter): cwt per acre</u>						
Mean	35.3	39.8	39.2		37.1	38.0	38.3
	<u>Kale, fresh weight: tons per acre</u>						
Mean	20.96	26.18	27.82		24.67	25.42	25.56

Mean dry matter % as harvested: Wheat 81.2
Barley 81.3
Oats 82.6

60/ca/2.3

	None	S1	Spray in 1959 S2	S3	A2	Mean
<u>Great Field I (R)</u>						
	<u>Sugar beet. Roots (washed): tons per acre</u>					
Mean	19.79	22.22	20.00	19.64	16.80	19.68
	<u>Sugar beet. Sugar percentage</u>					
Mean	17.0	16.4	16.2	16.3	16.9	16.5
	<u>Sugar beet. Total sugar: cwt per acre</u>					
Mean	67.2	72.6	64.9	63.9	57.0	65.0
<u>Great Knott I (R)</u>						
	<u>Wheat, grain (at 85% dry matter): cwt per acre</u>					
Mean	50.2	53.7	50.8	52.0	52.3	51.5
	<u>Barley, grain (at 85% dry matter): cwt per acre</u>					
Mean	36.1	37.6	38.2	37.6	40.0	37.6
	<u>Oats, grain (at 85% dry matter): cwt per acre</u>					
Mean	32.3	32.0	30.1	30.6	33.5	31.8
	<u>Kale, fresh weight: tons per acre</u>					
Mean	25.19	23.32	21.72	18.78	22.60	22.80
	<u>Sugar beet. Roots (washed): tons per acre</u>					
Mean	17.88	17.02	15.48	16.58	15.64	16.74
	<u>Sugar beet. Sugar percentage</u>					
Mean	16.9	16.7	16.7	16.6	16.6	16.7
	<u>Sugar beet. Total sugar: cwt per acre</u>					
Mean	60.3	56.8	51.8	55.2	51.8	56.0
Mean dry matter % as harvested:						
			Wheat	80.0		
			Barley	81.8		
			Oats	84.5		

60/Cd/2.4

<u>Broad Mead I (W)</u>						
	None	S1	Spray in 1959 S2	S3	A2	Mean
<u>Wheat, grain (at 85% dry matter): cwt per acre</u>						
Mean	36.6	38.2	36.7	37.6	40.5	37.9
<u>Kale, fresh weight: tons per acre</u>						
Mean	20.92	24.36	22.79	20.92	21.80	22.16
<u>Sugar beet. Roots (washed): tons per acre</u>						
Mean	15.83	17.80	13.75	16.27	16.32	15.99
<u>Sugar beet. Sugar percentage</u>						
Mean	14.9	15.6	14.2	15.1	15.2	15.0
<u>Sugar beet. Total sugar: cwt per acre</u>						
Mean	47.1	55.5	39.0	49.0	49.6	48.0
<u>Sugar beet. Tops: tons per acre</u>						
Mean	23.47	24.75	24.46	22.10	23.57	23.67

Mean dry matter % as harvested: Wheat 76.3

60/Ca/2.5

Great Hill (W)

	Spray and treatment in 1959							
	None	S1	S2	S3	S4	A2	M	Mean
<u>Barley, grain (at 85% dry matter): cwt per acre</u>								
Mean	22.4	23.4	24.2	21.4	20.3	18.0	20.2	21.4
<u>Oats, grain (at 85% dry matter): cwt per acre</u>								
Mean	10.9	10.4	11.2	8.9	7.2	8.4	6.9	9.1
<u>Sugar beet. Roots (washed): tons per acre</u>								
Mean	15.26	17.72	13.35	15.88	9.47	16.76	17.48	15.13
<u>Sugar beet. Sugar percentage</u>								
Mean	15.9	16.8	16.0	16.2	16.2	16.2	16.9	16.3
<u>Sugar beet. Total sugar: cwt per acre</u>								
Mean	48.6	59.4	42.6	51.4	30.7	54.2	59.2	49.4
<u>Sugar beet. Tops: tons per acre</u>								
Mean	14.00	13.65	13.48	17.39	8.87	13.39	14.35	13.59

Mean dry matter % as harvested: Barley 81.5
Oats 73.0

Sprays

S = Simazine
A = Atrazine

Levels

1 = 1 lb in 40 gallons per acre
2 = 2 lb in 80 gallons per acre
3 = 3 lb in 120 gallons per acre
4 = 4 lb in 160 gallons per acre

50% active
material

M = Normal mechanical weed control.