


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/R/CE/2 and 60/W/CE/2 Spring Beans - Control of Weeds (Spraying)

Rothamsted Research

Rothamsted Research (1961) *60/R/CE/2 and 60/W/CE/2 Spring Beans - Control of Weeds (Spraying)* ;
Yields Of The Field Experiments 1960, pp 115 - 116 - DOI:

<https://doi.org/10.23637/ERADOC-1-180>

60/Ce/2.1

SPRING BEANS

Control of weeds by triazine sprays - Rothamsted (R) Deacons Field and Woburn (W) Warren Field 1960.

Design: 4 randomised blocks of 6 plots each.

Area of each plot (acres):

Deacons Field (R): 0.0333. Area harvested: 0.0139.

Warren Field (W): 0.0303. Area harvested: 0.0126.

Treatments:

Simazine (2-chloro-4-6-bis-ethylamino-s-triazine - 50% active material):

Without inter row cultivations:

None; (S₀)

1 lb per acre; (S₁)

2 lb per acre; (S₂)

3 lb per acre; (S₃)

With normal inter row cultivations:

None; (MS₀)

2 lb per acre; (MS₂)

Basal dressing per acre: Deacons (R): 2 $\frac{3}{4}$ cwt compound fertiliser (12% P₂O₅, 24% K₂O) placement drilled. Warren Field (W): 3 $\frac{1}{2}$ cwt compound fertiliser (14% P₂O₅, 28% K₂O) placement drilled.

Cultivations, etc.:-

Deacons Field (R): Ploughed: Oct 13 - 26, 1959 and Mar 1 - 7, 1960.

Seed placement drilled at 200 lb per acre, with basal fertiliser:

Mar 19. Sprays applied to appropriate plots: Mar 25. Sprayed

with demeton-methyl at 12 fluid oz in 40 gallons per acre:

June 13. Combine harvested: Sept 8. Variety: Gartons Spring

Tick. Previous crop: Barley.

Warren Field (W): Ploughed: Oct 22 - 26, 1959. Ground chalk

applied at 18 - 20 cwt per acre: Mar 8 - 12, 1960. Seed

placement drilled at 200 lb per acre, with basal fertiliser:

Mar 24. Sprays applied to appropriate plots: Mar 25. Sprayed

with demeton-methyl at 12 fluid oz in 40 gallons per acre:

June 15. Combine harvested: Sept 21. Variety: Gartons Spring

Tick. Previous crop: Spring wheat.

Note: Weed counts were made in July.

Standard errors per plot, Grain (at 85% dry matter):

Deacons Field (R): 4.09 cwt per acre or 16.5% (15 d.f.)

Warren Field (W): 1.83 cwt per acre or 13.4% (15 d.f.)

60/0e/2.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

Treatment						Mean
S ₀	S ₁	S ₂	S ₃	MS ₀	MS ₂	
Deacons Field (R)						
25.0	23.6	26.2	26.6	23.8	23.3	24.7
(±2.04)						
Warren Field (W)						
11.3	13.0	12.1	12.6	17.0	16.0	13.6
(±0.91)						

Mean dry matter % as harvested:

Deacons Field (R): 73.9

Warren Field (W): 73.6

Treatments

Simazine without inter row cultivations

S₀ = None

S₁ = 1 lb per acre

S₂ = 2 lb per acre

S₃ = 3 lb per acre

Simazine with normal inter row cultivations

MS₀ = None

MS₂ = 2 lb per acre

Note. Deacons (R): It is suspected that certain plots in each block were affected by gross soil differences and this matter is under investigation.