


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/R/CG/1 Sugar Beet - Control of Virus Spread (Sprays)

Rothamsted Research

Rothamsted Research (1961) *60/R/CG/1 Sugar Beet - Control of Virus Spread (Sprays)* ; Yields Of The Field Experiments 1960, pp 126 - 127 - DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/Cg/1.1

SUGAR BEET

Control of virus spread by insecticide spray - Fosters 1960.

Design: 4 × 4 Latin square.

Area of each plot: 0.0168 acres. Area harvested: 0.0084 acres.

Treatments.

Unsprayed (0);

Sprayed when the crop was 0.3% infested with Myzus persicae
(May 23, 1960) (E);

Sprayed on receipt of spray warning on May 30, 1960 (2 plots per
row or column*) (N);

The spray used was demeton-methyl at 12 fluid oz in 60 gallons of
water per acre.

*Note. One treatment was to include a second spray, depending on the
development of a sufficiently large aphid population; however, the
population remained very small, and the spray was omitted.

Basal dressing per acre: 5 cwt salt, 6 cwt compound fertiliser (16% N,
9% P₂O₅, 9% K₂O).

Cultivations, etc.: Ploughed: Jan 7, 1960. Salt applied: Feb 24.
Basal compound fertiliser applied: Apr 2. Seed drilled at 19 lb
per acre: Apr 7. Singled: May 25. Lifted: Nov 8. Variety:
Klein E. Previous crop: Kale and mangolds in strips with small
plots of potatoes.

Note: Regular counts of aphids and virus yellows were made.

Standard error per plot.

Total sugar: 5.90 cwt per acre or 8.4% (7 d.f.)

60/Cg/1.2

Summary of Results

	None	Spray E	N	Mean
<u>Roots (washed): tons per acre</u>				
Mean	19.50	20.61	21.96	21.01
Increase		1.11	2.46	
<u>Sugar percentage</u>				
Mean	16.8	16.6	16.6	16.7
Increase		-0.2	-0.2	
<u>Total sugar: cwt per acre</u>				
Mean	65.4 (± 2.95)	68.6	73.2 (± 2.09)	70.1
Increase		3.2 (± 4.17)	7.8 (± 3.61)	

Sprays

O = Unsprayed

E = Sprayed when the crop was 0.3% infested with Myzus persicae

N = Sprayed on receipt of spray warning on May 30, 1960 (2 plots per row or column)