


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/W/B/7 Irrigation

Rothamsted Research

Rothamsted Research (1961) *60/W/B/7 Irrigation* ; Yields Of The Field Experiments 1960, pp 65 - 70
- DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/B/7.1

IRRIGATION EXPERIMENT

Revised 1960 (the 10th year)

The effects of irrigation and nitrogen - Woburn Butt Close 1960.

For details of previous cropping, treatments etc. see "Details of the Classical and Long Term Experiments" 1956.

The 3 course rotation is now as follows:-

1st year: early potatoes (following 1959 sugar beet).

2nd year: barley (following 1959 spring beans).

3rd year: winter beans (following 1959 spring wheat).

The fourth series carries a long term ryegrass ley for cutting.

Area of each whole plot (acres): Winter beans: 0.0555.

Sub plots (acres): Grass: 0.0264; remainder: 0.0277.

Area harvested (acres): Early potatoes: 0.0075; barley: 0.0110;
winter beans: 0.0177; grass: 0.0165.

Design: 4 series (1 in each crop) each containing 12 whole plots.

The bean plots are no longer split for the application of dung.

Treatments.

Early potatoes: all combinations of:-

Whole plots: Irrigation: None (0); full (C).

Weed control: normal cultivations after planting (no spray);
simazine spray at 2 lb in 40 gallons per acre (used as
pre-emergence weedkiller) without cultivations.

Sub plots: Nitrogen: None, 0.6 cwt N per acre as sulphate of
ammonia.

Note: After the early potatoes are lifted certain plots are sown
with trefoil as a green manure for barley. Provision is made
for a comparison of none v. irrigation applied to the trefoil.
No irrigation was applied in 1960.

Barley: All combinations of:-

Whole plots: Irrigation: None (0), full (C).

Sub plots: Nitrogen*: None, 0.2 cwt N per acre as 'Nitro-Chalk'.

Winter beans:

Whole plots: Irrigation: None (0), three systems (A, B, C -
see below).

Grass: all combinations of:-

Whole plots: Irrigation: None (0), full (C).

Potash: None, 0.6 cwt K_2O per acre as muriate of potash
applied after the 1st cut and once again in mid-season.

Sub plots: Nitrogen*: None, 0.3 cwt N per acre as 'Nitro-Chalk'
in spring and after each cut except the last.

*Note: In addition to basal dressing.

60/B/7.2

Rainfall and Irrigation: inches

Week ending	Rain-fall	Grass	Barley	Potatoes	Beans		
		C	C	C	A	B	C
May 2	0.13	0.50	0.50	0.50	-	0.50	0.50
9	0.02	0.50	-	-	-	0.50	0.50
16	0.89	0.62	-	-	-	0.50	0.50
23	0.23	-	0.50	-	-	0.50	0.50
30	0.01	0.50	0.50	0.50	-	0.67	0.67
June 6	-	0.67	0.50	0.50	0.67	0.67	0.67
13	1.26	-	-	-	-	-	-
20	0.68	-	-	-	0.20	-	0.20
27	1.68	0.50	-	0.75	-	-	-
July 4	0.01	-	-	-	-	-	-
11	1.23	-	-	-	-	-	-
18	1.19	-	-	-	-	-	-
25	0.37	-	-	-	-	-	-
Aug 1	0.38	-	-	-	-	-	-
8	0.23	-	-	-	-	-	-
15	0.88	-	-	-	-	-	-
22	0.16	-	-	-	-	-	-
29	0.87	0.50	-	-	-	-	-
Sept 5	1.60	-	-	-	-	-	-
12	0.16	-	-	-	-	-	-
19	1.52	-	-	-	-	-	-
26	(0.63)	-	-	-	-	-	-
Oct 3	(0.64)	-	-	-	-	-	-
Total	14.77	3.79	2.00	2.25	0.87	3.34	3.54

Basal dressings (per acre):

Early potatoes: 0.60 cwt N as sulphate of ammonia; 0.75 cwt P_2O_5 and 1.50 cwt K_2O as compound fertiliser (14% P_2O_5 , 28% K_2O).

Barley: 0.2 cwt N, 0.2 cwt P_2O_5 and 0.3 cwt K_2O as compound fertiliser (12% N, 12% P_2O_5 , 18% K_2O).

Winter beans: 0.3 cwt P_2O_5 , 0.6 cwt K_2O placement drilled as compound fertiliser (10% P_2O_5 , 20% K_2O).

Grass: 0.3 cwt N as 'Nitro-Chalk' in spring and again after each cut except the last, and 0.6 cwt P_2O_5 and 1.2 cwt K_2O as compound fertiliser (14% P_2O_5 , 28% K_2O).

Cultivations, etc.:

Early potatoes: Ploughed: Nov 20, 1959. PK compound applied: Apr 4, 1960. Sulphate of ammonia applied: Apr 6. Machine planted: Apr 7. Appropriate plots sprayed with simazine: Apr 15. Earthed up (except simazine plots): June 4. Haulm destroyed mechanically: July 13. Lifted: July 15. Trefoil sown at 30 lb per acre: July 21. Variety: Arran Pilot.

60/B/7.3

Barley: Ground chalk applied at 3 tons per acre: Sept 8, 1959.

Ploughed: Sept 9 and Nov 21. Seed drilled at $2\frac{1}{4}$ bushels per acre: Mar 19, 1960. Fertilisers applied: Mar 21. Sprayed with DNBP at 10 pints in 80 gallons per acre: May 16. Combine harvested: Aug 13. Variety: Proctor.

Winter beans: Ploughed: Sept 7, 1959. Seed placement drilled at 275 lb per acre with PK compound: Nov 5. Harvested: Aug 10 and Aug 26, 1960. Variety: Rothamsted S.Q.

Grass: Ground chalk applied at 18 cwt per acre: Sept 23, 1959. Seed sown at 30 lb per acre: Oct 20. 'Nitro-Chalk' and PK compound applied: Apr 1, 1960. Cut 8 times (all plots): May 10, May 31, June 22, July 18, Aug 8, Aug 30, Sept 23, Nov 8. 'Nitro-Chalk' applied after each cut except the last. Muriate of potash applied to appropriate plots after 1st and 4th cuts. Variety: S22 Italian ryegrass.

Standard errors per plot.

Early potatoes. Total tubers

Whole plot: 0.708 tons per acre or 7.2% (4 d.f.)

Sub plot: 0.630 tons per acre or 6.4% (8 d.f.)

Barley, (grain at 85% dry matter)

Whole plot: 2.51 cwt per acre or 10.2% (8 d.f.)

Sub plot: 2.25 cwt per acre or 9.1% (10 d.f.)

Winter bean, (grain at 85% dry matter)

Whole plot: 2.80 cwt per acre or 9.5% (6 d.f.)

Cut grass, dry matter

1st cut:

Whole plot: 0.58 cwt per acre or 7.3% (6 d.f.)

Sub plot: 1.44 cwt per acre or 18.2% (8 d.f.)

Total of cuts 2-4

Whole plot: 2.03 cwt per acre or 4.2% (6 d.f.)

Sub plot: 1.98 cwt per acre or 4.1% (8 d.f.)

Total of cuts 5-8

Whole plot: 1.48 cwt per acre or 3.7% (6 d.f.)

Sub plot: 2.59 cwt per acre or 6.6% (8 d.f.)

Total of all 8 cuts

Whole plot: 2.55 cwt per acre or 2.7% (6 d.f.)

Sub plot: 3.96 cwt per acre or 4.1% (8 d.f.)

Early potatoes, Total tubers: tons per acre

Barley, (Grain at 85% dry matter): cwt per acre

Winter beans, (Grain at 85% dry matter): cwt per acre

*** For use in horizontal and diagonal comparisons only.

60/B/7.5

Cut grass, Dry matter: cwt per acre

<u>1st cut</u>					
K ₂ O: cwt per acre (including basal) in 1959		Irrigation 0 C			
		(±0.33)			
1.2		7.4	7.5	K ₂ O: cwt per acre including basal	
2.4		4.8	11.8		
N: cwt per acre				1.2 2.4	Mean
		(±0.48)*		(±0.48)*	
0.3		5.8	9.5	8.0	7.7
0.6		6.4	9.7	6.9	8.0
Mean	(±0.24)	6.1	9.6	7.4	7.9
Difference	(±0.83)	+0.6	+0.2	-1.1	+0.3 (±0.59)

Total of cuts 2 - 4

K ₂ O: cwt per acre including basal		Irrigation 0 C			
		(±1.17)			
1.2		41.4	51.5	K ₂ O: cwt per acre including basal	
1.8		44.5	55.7		
N: cwt per acre				1.2 4.8	Mean
		(±1.01)*		(±1.01)*	
0.3		41.7	48.7	44.2	45.2
0.6		44.1	58.4	48.6	51.3
Mean	(±0.83)	42.9	53.6	46.4	48.3
Difference	(±1.14)	2.4	9.7	4.4	6.1 (±0.81)

*For use in horizontal and diagonal comparisons only.

Mean dry matter % as cut:

1st cut: 20.9

Total of cuts 2 - 4: 19.0

60/B/7.6

Cut grass, Dry matter: cwt per acreTotal of cuts 5 - 8

K ₂ O: cwt per acre including basal	Irrigation		K ₂ O: cwt per acre including basal		Mean
	0	C			
	(±0.85)				
1.2	38.9	35.4			
2.4	40.8	43.1			
N: cwt per acre			1.2	2.4	
	(±0.96)*		(±0.96)*		
0.3	36.7	36.9	35.0	38.7	36.8
0.6	43.0	41.5	39.3	45.2	42.3
Mean (±0.60)	39.9	39.2	37.2	41.9	39.5
Difference (±1.50)	+6.3	+4.6	+4.3	+6.5	+5.5 (±1.06)

Total of cuts 1 - 8

K ₂ O: cwt per acre including basal	Irrigation		K ₂ O: cwt per acre including basal		Mean
	0	C			
	(±1.47)				
1.2	87.7	94.3			
2.4	90.1	110.6			
N: cwt per acre			1.2	2.4	
	(±1.55)*		(±1.55)*		
0.3	84.2	95.2	87.1	92.3	89.7
0.6	93.5	109.7	94.9	108.3	101.6
Mean (±1.04)	88.9	102.5	91.0	100.3	95.7
Difference (±2.29)	9.3	14.5	7.8	16.0	11.9 (±1.62)

*For use in horizontal and diagonal comparisons only.

Mean dry matter % as cut:

Total of cuts 5 - 8: 15.0

Total of cuts 1 - 8: 17.3