Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1907



Full Table of Content

Annual Supplement for 1907

Annual Supplement for 1907, Rothamsted Research (1908) Yields Of The Field Experiments 1907, pp 1 - 7 - DOI: https://doi.org/10.23637/ERADOC-1-144

LAWES AGRICULTURAL TRUST.

Rothamsted Experimental Station, Harpenden.

Annual Supplement for 1908

to "Guide to the Experimental Plots, 1906,"

CONTAINING THE YIELDS PER ACRE FOR 1907.

In every case the page, table, and plot numbers refer to the "Guide," it being understood that no change is made in the manuring, etc., there described.

A. D. HALL. DIRECTOR.

ST. ALBANS:

PRINTED BY W. CARTMEL AND SONS, 6, VICTORIA STREET.
1908.

METEOROLOGICAL RECORDS, 1907.

(See "Guide," page 16, Table IX.)

	Y	Rain.		Drai	Drainage through				
	Total Fall.		No. of Rainy Days.		soil.		Bright	Tempe	rature
		Days.				Sun- shine.			
	5-inch Funnel Gauge.	Toooth Acre Gauge.	Acre Gauge.	20 ins. deep.	40 ins. deep.	60 ins. deep.		Max.	Min.
January	Inches.	Inches.	No. 10	Inches. 0.863	Inches. 0.956	Inches.	Hours.	°F.	°F.
February	1.452	1.476	15	0.958	1.047	1.025	85.2	41.8	30-7
March	1.285	1.312	13	0.057	0.155	0.107	206.4	52.9	32.2
April	2.855	2.846	17	0.567	0.539	0.509	143.2	53.9	37.7
May	2.377	2.396	18	0.288	0.384	0.356	164-4	59.5	42.9
June	2.484	2.609	20	0.358	0.391	0.366	160.1	62.5	48.4
July	2.165	2.209	16	0.276	0.259	0.236	170.6	65.1	48.9
August	1.682	1.802	14	0.003	0.016	0.013	174.5	66.6	50.2
September	0.721	0.778	8	***	**	***	185.1	65.9	46.8
October	4.876	4.890	23	2.943	2.830	2.713	97.3	56.1	42.2
November	2-424	2.439	18	1.910	1.982	1.940	58.0	49.4	37.8
December	3.435	3.396	20	3.064	3.201	3.165	45.8	45.0	34.9
Total or Mean	26.992	27-407	192	11.287	11.760	11:396	1557-1	55.1	40.4

MANGEL WURZEL. BARN FIELD, 1907.

(See "Guide," page 11, Table VI.)

						C	ross-dressings	•	
2.0000	Strip	0.			N.		A.	A C.	C.
Strip.	Manures.	N	lone.	N	itrate Soda.	of	Ammonium Salts.	Rape-cake & Ammonium Salts.	Rape Cake.
1	Dung only	βR. γL.	26.00 3.64		Tons. 41.42 4.64		Tons. 33-52 5-27	Tons. 34·29 4·90	Tons. 35.02 5.17
2	Dung, Super., Potash	(R. L.	26·52 3·33		42·13 4·61		41.68 6.64	43·52 7·08	40.74 5.34
4	Complete Minerals	∫R. (L.	5•95 1•09	alera de	30·46 35·15 4·00 4·94	とうとう	26.68 3.42	40-97 5-25	33·09 4·11
5	Superphosphate only	(R. (L.	6·21 1·17		24·62 3·42		10.88 2.86	11.26 2.18	15·43 2·18
6	Super. and Potash	(R. (L.	5·78 1·05		25.05 3.14		25:22 3:44	35·88 5·68	28·15 2·84
7	Super., Sulph. Mag. & Chloride Sodium	(R. (L.	6·59 1·27		26·54 3·75		26·52 3·44	34·38 5·29	30-59 3-76
8	None	(R.	5·15 1·06	1	18.60 3.84		9·87 3·03	10·90 2·26	13.24 2.40

HAY. THE PARK GRASS PLOTS, 1907.

(See "Guide," page 19. Table XI.)

Plot.	Manuring.	Yield of Hay per acre.				
. 101.	Manuring,	1st Crop.	2nd Crop.	Total.		
POPULATION AND ADDRESS OF THE PARTY.		Cwt.	Cwt.	Cwt.		
3 (12 (Unmanured	21.3	1.8	23.1		
12 (25·3 26·1	4·3 2·2	29·6 28·3		
2	Unmanured (1) Ammonium salts alone (1)	34.4	5.3	39.7		
4-1	Ammonium salts alone (1) Superphosphate of Lime	25.0	2.0	27.0		
	Mineral Manure without Potash	36.3	5.9	42.2		
8 7 6	Complete Mineral Manure	57.2	13.2	70.4		
6	As 7, 1869 and since (2)	49.6	13.0	62.6		
15	As 7, 1876 and since (3)	46.0	11.7	57.7		
5	Superphosphate and Potash, 1898					
	and since	24.5	2.1	26.6		
17	Nitrate of Soda alone	38.8	7.9	46.7		
4-2	Superphosphate and Ammsalts	40-6	1.9	42.5		
10	Mineral Manure (without Potash) and	F0.0	4.0	***		
	Ammsalts	50.0	4.7	54.7		
9	Complete Mineral Manure and Amm salts	65:0	7.9	72.9		
13	Dung and Fish Guano, once in 4 yrs.	52.2	15.5	67.7		
11-1	Complete Mineral Manure and extra	02.2	200	0		
	Ammsalts	46.7	24.0	70.7		
11-2	As 11-1, and Silicate Soda	68.3	29-2	97.5		
16	Complete Mineral Manure and Nit.					
9550	Soda=43 lb. N	50.4	12.2	62.6		
14	Do. do. do. and Nit.	0.0000		27.2		
10000	Soda=86 lb. N	51.2	13.3	64.5		

Quick Lime (ground), at the rate of 2000 lb. per acre, applied to the South half of plots 1 to 4-2, 7 to 11-2, 13 and 16, in January, 1907.

Received Farmyard Dung, 8 yrs., 1856-63.
 Ammonium salts alone, previous to 1869.

(3) Nitrate of Soda alone previously.

BOTANICAL COMPOSITION, PER CENT.

First Crop, 1907.

(See "Guide," page 20, Table XII.)

Plot.	Manuring.		Gramineæ.	Leguminosæ.	Other Orders
			Per cent.	Per cent.	Per cent.
3	Unmanured		51.6	6.2	42.2
4-1	Superphosphate of Lime		54.4	5.2	40.4
8	Mineral Manure without Pota	sh	44.8	15.2	40.0
7	Complete Mineral Manure		53.3	29.3	17-4
6	As 7, 1869 and since (2)		44.9	38-7	16-4
15	As 7, 1876 and since (3)		37.8	49-9	12.3

WHEAT. BROADBALK FIELD, 1907.

(See " Guide," page 26, Table XIV.)

		Dresse	l Grain.	
Plot.	Manuring.	Yield.	Weight per Bushel.	Straw.
		Bushels.	lbs.	Cwt.
2	Farmyard Manure	33.7	60·5 60·7	56.6
3 5 6 7 8 9	Unmanured	9·1 11·5	60-6	9·8 15·1
o o	Complete Mineral Manure	23.9	60.4	31.4
7	As 5, and single Ammsalts As 5, and double do	33.6	60.5	55.8
8	As 5, and double do As 5, and treble do	34.7	59.4	71.6
9	As 5, and single Nitrate Soda	30.2	60.5	46.0
10	Double Ammsalts alone	27.6	60.2	33.1
11	As 10, and Superphosphate	32.1	60.7	40.5
12	,, and Super and Sulph. Soda	41.6	60-9	51.3
13	,, and Super and Sulph. Potash	34.7	60-2	57.3
14	,, and Super and Sulph. Mag.	36.6	60-9	46.9
15	Double Ammsalts in Autumn, and		1	
	Minerals	33.1	61-6	48.6
16	Double Nitrate and Minerals	34.7	60.4	65.7
17	Minerals alone, or Double Amm	*10.9	61.3	13.0
18	salts alone, in alternate years	+31.0	60-7	44.3
19	Rape Cake alone	29.0	60-3	39.3

BARLEY. HOOS FIELD, 1907.

(See " Guide," page 33, Table XVI.)

		Dressed	l Grain.	
Plot.	Manuring.	Yield.	Weight per Bushel.	Straw
10	No Minerals, and no Nitrogen	Bushels.	lbs. 53·5	Cwt.
20	Superphosphate only	13.0	56.5	9.2
30	Alkali salts only	8.1	54.3	14.8
4 0	Complete Minerals	13.2	55.0	24.3
1 A	Ammsalts only	19.9	53.6	14.2
2 A	Superphos. and Ammsalts	28.4	55.2	20.0
3 A	Alkali salts and Ammsalts	20.1	54.4	15.8
4 A	Complete Minerals and Ammsalts.	32.7	55.4	25.5
1 N	Nitrate of Soda alone	22.3	54.0	17.2
2 N	Superphos. and Nitrate Soda	29.0	56.1	25.4
3 N	Alkali salts and Nitrate Soda	21.2	55.0	15.9
4 N	Complete Minerals and Nitrate Soda	31.7	56.1	27.2
1 C	Rape Cake alone	29.1	55-3	21.2
2 C	Superphos. and Rape Cake	28.8	56.1	21.6
3 C	Alkali salts and Rape Cake	26.3	55.9	19-9
4 C	Complete Minerals and Rape Cake	31.1	56.7	23.4
7-1	Unmanured (after Dung, 1852-71)	15.5	55.3	18.4
7-2	Farmyard Dung	42.1	57.1	38.0

^{*} Produce by Minerals. † Produce by Ammonium-salts.

BARLEY. HOOS FIELD, 1907.

(Previous cropping: Potatoes, 1876-1901; Barley, 1902 and 1903; Oats, 1904; Barley, 1905 and 1906).

(See "Guide," page 40, Table XIX.)

	Manures applied	Dressed	Grain.		Total Produce	
Plot.	to the Potatoes, 1876-1901. Unmanured since.	Yield.	Weight per Bushel.	Straw.		
		Bushels.	lbs.	Cwt.	lbs.	
1	Unmanured	6-2	56.0	4.2	838	
2	Unmanured 1882 and since, previously Dung alone	10-3	57-3	7-0	1400	
3	Dung 1883-1901	18.9	56.7	15.4	2855	
4	Dung 1883-1901	19-1	56.8	14.9	2818	

WHEAT AFTER FALLOW (without manure 1851 and since).

HOOS FIELD, 1907.

(See "Guide," page 41, Table 20).

Dressed C	rain!	 	Yield—14·3 bushels. Weight per bushel—58·6 lbs.
Straw		 	19·5 cwt.
Total Pro		 	3094 lbs.

INOCULATION OF LEGUMINOUS PLANTS. HOOS FIELD.

(See "Guide" page 40, and plan page 37).

PRODUCE OF RED CLOVER (HAY) IN 1907.

1. EFFECT OF INOCULATING THE SOIL.

Plot.	Soil inoculated with-	Mean of Plots 6, 8, and 10 1st and 2nd Crops
Α	Hiltner's Preparation from Munich	Cwt. 66·1
В	Moore's Preparation from the United States	57.4
С	Soil from a field which had carried Red Clover in 1904	59-0
D	Left uninoculated	56-6

2. EFFECT OF PAST MANURING.

Plot.		Mean of Plots, A, B. C, D, 1st and 2nd Crops.
6	Nitrate of Soda 1876-1901, since unmanured	Cwt. 58·7
8	Nitrate of Soda and Mixed Minerals 1876-1901, since unmanured	64.5
10	Mixed Minerals only 1876-1901. since unmanued	56.2

3. DETAILS OF THE ABOVE.

Plot.	1st Crop.	2nd Crop.	Total.
25 31	Cwt.	Cwt.	Cwt.
6 A	42.0	20.6	62.6
6 B	40-5	18-4	58.9
6 C	40.5	18-1	58.6
6 D	36-0	18.9	54.9
	45.0	07.0	50.0
8 A	45.0	27.0	72.0
8 B	39.0	21.5	60.5
8 C	40.5	23.1	63.6
8 D	40.5	21.3	61.8
10 A	37.5	26.1	63.6
10 B	33.0	19-8	52.8
10 C	34.5	20.5	55.0
10 D	33.0	20.3	53.3

LITTLE HOOS FIELD, 1904-07.

RESIDUAL VALUE OF VARIOUS MANURES. (See "Guide," pages 41 and 42).

Total Produce—Grain and Straw or Roots and Leaves, per acre.

Series and Plot.	Manuring				Swedes 1904.	Barley 1905.	Mangels 1906.	Spring Whea 1907.
A 1	Unmanured				Tons. 10·3	1bs. 2323	Tons. 17·1	lbs. 3650
3	Dung (ordinary), 1904 o	000	***	***	13.1	4649	18-2	4673
4	,, ,, 1905 ,, 1906	**		***	8.8	3501 2269	17·5 18·2	5393 5471
5	., ., 1907	"			9.8	2402	14.9	6903
B 1	Dung (cake-fed), 1904 c	nly			15.7	4177	19.4	4319
2 3	Unmanured	***	***		10.0	2417	16.2	4025
4	Dung (cake-fed), 1905 o		***	***	9·5 11·4	5530 2772	18·5 25·6	5497 6489
5	,, ,, 1906	"			9.4	2649	14.4	9407
C 1	Shoddy, 1904 only				14.7	3656	21.0	4667
2	,, 1905 ,,				11.1	4363	23.6	4550
3	Unmanured	***		***	10.6	2588	17.7	4334
4 5	Shoddy, 1906 only ,, 1907 ,,				10·7 10·3	$\frac{2512}{2615}$	24·2 16·9	6231 7495
D 1	Guano, 1904 only		-			0550	00.7	10-0
2	, 1905 ,,	***	***		14·6 11·0	2550 5176	20·1 19·7	4056 4165
3	,, 1906 ,,		100		10.9	2857	25.6	4846
4	Unmanured	***			10.6	2985	18.7	4618
5	Guano, 1907 only	•••	***		10.6	2680	17.4	7375
E 1	Rape-cake, 1904 only				14-1	2674	17.8	3887
2 3	,, ,, 1905 ,,	***	**		11.2	4185	17.9	4326
4	,, ,, 1906 ,, ,, 1907 ,,		***		9·5 10·5	2645 2734	22·7 19·4	4584 6619
5	Unmanured				10.8	2769	19.5	4527
F 1	Unmanured				11-7	3132	22.9	4749
2	Superphosphate, 1904 o	nly			12.2	3025	23.2	5064
3		**	***		10.2	3949	23.6	4956
4 5	1007	"			9.7	3913 4221	24·1 23·6	5419 5698
G 1	Bone Meal, 1904 only				12-9	3176	23.1	5000
2	., ., 1905			:::	10.1	3636	22.1	5203 5821
3	Unmanured		***	***	10.2	3495	20.6	5491
5	Bone Meal, 1906 only	***	•••		9.9	3450	22.6	6043
ð	,, ,, 1907 ,,				9-2	3525	22.1	6276
H 1	*****				11.8	4400	20-5	6285
3	,, ,, 1905 ,, ,, ,, 1906 ,,		***	***	9-4	4002	21.3	5930
4	1007	•••			9.1	3662 3624	21·4 17·0	5860 5816
5	Unmanured				8.6	3293	17.4	5933