


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1974

[Full Table of Content](#)



74/R/RN/9 Cereal Disease Reference Plots - Beans, Oats, Wheat

74/R/RN/9 Cereal Disease Reference Plots - Beans, Oats, Wheat, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 104 - 105 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/R/RN/9

CEREAL DISEASE REFERENCE PLOTS

Object: To study the effects of intensive cereal cropping on the incidence of soil-borne diseases, especially in relation to seasonal variation - Pennell's Piece.

Sponsors: D.B. Slope, E.W. Broom, G.A. Salt.

The twelfth year, winter wheat, spring oats, beans.

For previous years see 63/C/10(t), 64-65/C/9, 66/C/7, 67-68/C/5 and 69-73/R/RN/9.

Design: 2 randomised blocks of 6 plots.

Whole plot dimensions: 17.1 x 4.27. Area harvested: 0.00479.

Treatments: Previous crops (1963-73):-

PREVCROP

1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973

O	W	W	W	BE	O	W	W	W	BE	O	W/W/BE/O
W	W	W	BE	O	W	W	W	BE	O	W	W/BE/O/W
W	W	BE	O	W	W	W	BE	O	W	W	BE/O/W/W
W	BE	O	W	W	W	BE	O	W	W	W	-
BE	O	W	W	W	BE	O	W	W	W	BE	-
W	W	W	W	W	W	W	W	W	W	W	W/W/W/W

where: W = wheat, BE = beans, O = oats

Standard applications:

Winter wheat: Manures: (0:14:28) at 270 kg, combine drilled, 'Nitro-Chalk' at 500 kg. Weedkillers: Chlortoluron ('Dicurane 80 WP' at 4.5 kg in 220 l), dicamba with mecoprop and MCPA ('Banlene Plus' at 5.6 l in 370 l).

Spring beans: Manures: (0:14:28) at 450 kg, placement drilled.

Insecticide: Demeton-s-methyl at 0.25 kg in 220 l.

Spring oats: Manures: (0:14:28) at 270 kg, combine drilled, 'Nitro-Chalk' at 200 kg. Weedkiller: Dicamba with mecoprop and MCPA ('Tetralex-Plus' at 7.0 l in 220 l).

Seed: Winter wheat: Cappelle, sown at 200 kg.

Spring beans: Minor, sown at 220 kg.

Spring oats: Manod, sown at 190 kg.

74/R/RN/9

Cultivations, etc.:— All plots: Ploughed: 17 Sept, 1973.

Winter wheat: Power harrowed, seed sown: 15 Oct. 'Dicurane' applied: 22 Oct. N applied: 22 Mar, 1974. 'Banlene Plus' applied: 6 May. Combine harvested: 27 Aug.

Spring beans: Power harrowed: 27 Mar, 1974. Seed sown and spring-tine cultivated: 27 Mar. Insecticide applied: 13 June. Combine harvested: 14 Oct.

Spring oats: Power harrowed: 27 Mar, 1974. Seed sown: 28 Mar. N applied: 8 Apr. Weedkiller applied: 28 May. Combine harvested: 27 Aug.

- NOTES: (1) Yields were taken for winter wheat only.
(2) Estimates were made in July of incidence of take-all (*Gaeumannomyces graminis*) and eyespot (*Cercospora herpotrichoides*) in the wheat.

TABLES OF MEANS

WINTER WHEAT

GRAIN: TONNES/HECTARE

PREVCROP				Mean
W/W/BE/O	W/BE/O/W	BE/O/W/W	W/W/W/W	
6.28	5.70	5.79	5.67	5.86

Mean D.M.% 83.6