

Broadbalk Beans

Number of samples analysed for crop nutrient content

Harvest year	Section	%N		%P		%K		%Ca		%Mg		%Na		%S	
		Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw
1968	2	19	19	19	19	19	19	19	19	19	19	19	19	NA	NA
1969	7	19	19	19	19	19	19	19	19	19	19	19	19	NA	NA
1970	4	19	19	19	19	19	19	19	19	19	19	19	19	NA	NA
1971	2	19	NA	19	NA	19	NA	19	NA	19	NA	19	NA	NA	NA
1972	7	19	19	19	19	19	19	19	19	19	19	19	19	NA	NA
1973	4	19	19	19	19	19	19	19	19	19	19	19	19	NA	NA
1974	2	19	19	19	19	19	19	19	19	19	19	19	19	NA	NA
1975	7	19	19	19	19	19	19	19	19	19	19	19	19	NA	NA
1976	4	19	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1977	2	19	19	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
1978	7	19	19	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

NA Not analysed

Field beans grown in one rotational section of Broadbalk every year 1968-1978 (Sections 2,4 and 7).
Grown in a three year rotation: wheat > potato > field beans

19 samples = sampled from all plots in that Section