

Hoosfield spring barley experiment plan and fertilizer treatments 1852-1967

DOI: 10.23637/rhb2-plan1852-1967-01

Prepared by: M.J. Glendining, CAS Department, Rothamsted Research, Harpenden, Herts, AL5 2JQ, UK.

Published by: Rothamsted Research

Date: April 2012

Description: Standardised experiment plan for the Hoosfield spring barley experiment, 1852-1967, with details of fertilizer and manure treatments (not to scale).

- Page 1: Cover page
- Pages 2-3: Experiment plan and fertilizer treatments, 1852-1967

Site: R/HB/2. Hoosfield, Rothamsted Experimental Farm, Rothamsted Research, West Common, Harpenden, Hertfordshire, AL5 2JQ, UK. Geolocation: 51.81206, -0.37608

Related Resources:

- Rothamsted Experimental Station (1970): *Details of the Classical and Long-term Experiments up to 1967,* pp18-21. <u>https://doi.org/10.23637/ERADOC-1-192</u>
- Rothamsted Research (2015) Hoosfield spring barley experiment plans and fertilizer treatments, 1968-2000. Electronic Rothamsted Archive, Rothamsted Research. <u>10.23637/rhb2-plans1968-2000-01</u>
- Rothamsted Research (2009) Hoosfield spring barley experiment plan and fertilizer treatments, 2001 onwards. Electronic Rothamsted Archive, Rothamsted Research. 10.23637/rhb2-plan2001-01

Cite as: Rothamsted Research (2012) *Hoosfield spring barley experiment plan and fertilizer treatments, 1852-1967. Electronic Rothamsted Archive, Rothamsted Research.* <u>10.23637/rhb2-</u> <u>plan1852-1967-01</u>

Funding: Rothamsted Research receives strategic funding from the UK Biotechnology and Biological Sciences Research Council (BBSRC). The Rothamsted Long-term Experiments National Capability is supported by the BBSRC Grant BBS/E/C/000J0300 and the Lawes Agricultural Trust.


Licence and conditions of re-use:



These plans are published under <u>the Creative Commons Attribution 4.0 International</u> licence. CC BY 4.00

You are free to adapt, copy, redistribute these plans but must provide appropriate credit using the provided citation, including the DOI and indicate any changes made. You must not apply additional restrictions on the licence.

Hoosfield spring barley 1852 - 1967



Hoosfield treatments 1852-1967

Fertilizer treatments (per hectare per year unless indicated)

Nitrogen (applied in the spring)

Series O	No N fertilizer
Series A	48kg N as ammonium sulphate
Series AA	48kg N as sodium nitrate
Series C	48kg N as castor bean meal since 1941; rape cake 1852-1940
N*	48kg N as sodium nitrate
N ⁺	48kg N as ammonium sulphate

Series treatments applied to strips 1-4 only. Not applied to strips 6 and 7

Organics (applied in the spring)

FYM 1852 -	35t Farmyard manure since 1852
(FYM) 1852-71	35t Farmyard manure, 1852-1871 only

Minerals (applied in the spring)

Р	35kg P as superphosphate
K	90kg K as potassium sulphate
Na	15kg Na as sodium sulphate
Mg	11kg Mg as magnesium sulphate
Si	450kg sodium silicate since 1868;
	450kg calcium silicate and sodium silicate 1862-67
(Ashes)	Ashes were added to minerals to aid spreading, 1852-1932 only

Lime:

Lime has been applied as required since the 1950s to maintain soil pH at a level that does not compromise yield. For more details see 'Details up to 1967'.

Notes:

A combined yield is given for plots 7-1 and 7-2 1852-1871. From 1872 onwards, plots 7-1 and 7-2 are harvested separately.

Cropping:

Spring barley grown every year except 1912, 1933, 1943 and 1967 when the whole experiment was fallowed to control weeds.

Reference:

Rothamsted Experimental Station (1970). *Details of the Classical and Long-term experiments up to 1967.* Lawes Agricultural Trust, Harpenden, UK https://doi.org/10.23637/ERADOC-1-192